

## Original Article

# Assessing the ecological quality status of arid mangroves in the Gulf of Oman, Iran, using benthic indices of AMBI, M-AMBI, and BENTIX

Mohadeseh Miri<sup>1</sup>, Jafar Seyfabadi<sup>1\*</sup>, Mehdi Ghodrati Shojaei<sup>1</sup>, Hassan Rahimian<sup>2</sup>

<sup>1</sup>Department of Marine Biology, Faculty of Natural Resources and Marine Sciences, Tarbiat Modares University, Noor, Iran.

<sup>2</sup>School of Biology, College of Science, University of Tehran, Tehran, Iran.

**Abstract:** Polychaetes are suitable indicators to evaluate the benthic ecological status and respond to natural and anthropogenic. We evaluated the ecosystem health of the mangroves of Azini and Gwadar based on benthic indices including AMBI, M-AMBI, and BENTIX using polychaete communities. The results showed that in both regions, EcoQ classifications ranged from "high" to "moderate" in BENTIX, "good" to "excellent" in AMBI, and "good" in M-AMBI. The M-AMBI was significantly correlated with sediment variables, including total organic matter (TOM), total organic carbon (TOC), and silt/clay. The result revealed a significant correlation between the biotic indices and the TOC content of sediments. According to the results, TOC can be used as a descriptor and indicator to evaluate the health of mangrove ecosystems in relation to benthic indices. In addition, it is necessary to combine several indices to assess the status of ecosystems.

*Article history:*

Received 2 January 2023

Accepted 25 April 2023

Available online 25 April 2023

*Keywords:*

Mangrove

Bioindicator

Ecological quality status

Benthic communities

## Introduction

The mangrove forests grow in estuaries and intertidal zones of the tropics and subtropics (Field et al. 1998). The mangrove ecosystem is a biologically active ecosystem with numerous ecological functions. Mangrove forests provide many benefits, including breeding grounds for fish and shellfish, birds, and other wildlife, preventing shoreline erosion, and protection during hurricanes and tidal waves (Quarto, 2005). In addition, they are responsible for providing many ecosystem goods and services, including natural barriers, carbon sequestration, and biodiversity (Duke et al. 2007). However, it has been threatened by urbanization, pollution, and overexploitation over the past decades (Alongi, 2002).

The spatial distribution patterns of polychaetes have been extensively investigated with respect to environmental variables since they constitute a dominant element of benthic communities (Tyler and Kowalewski, 2018). The abundance of polychaetes on estuarine macrofauna and their

occurrence under a variety of environmental conditions make them an excellent biological model for studying estuarine ecosystems (Schüller et al., 2009). The assemblages of polychaetes display changes in standing stock as a response to variables in the environment, with the increase in organic sediment content being one of the most significant impacts of anthropogenic activity (Dauvin et al., 2016; Alvarez-Aguilar et al., 2017). Polychaetes can be used as a bioindicator of organic pollution due to their high diversity, abundance, and functional significance. They are frequently used as ecological groups to determine ecosystem quality as part of biotic indices (Borja et al., 2014).

Several indices based on benthic invertebrate communities have been proposed to evaluate environmental health status, including the AZTI Marine Biotic Index (AMBI) (Borja et al. 2000), the multivariate AMBI (M-AMBI) (Muxika et al., 2007), and the BENTIX Index (Simboura and Zenetos 2002). AMBI is based on species assigned to one of five levels of sensitivity, ranging from very

\*Correspondence: Jafar Seyfabadi  
E-mail: jseyfabadi@gmail.com




Figure 1. Positions of the sampling sites of polychaetes in the Hara Biosphere Reserve, Gulf of Oman (the Gwadar and Azini estuaries).

sensitive to opportunistic species (Borja et al., 2000). M-AMBI is a multivariate analysis that combines Shannon-Wiener diversity, species richness, and AMBI (Muxika et al., 2007). Benthic index calculates the relative contributions of tolerant and sensitive taxa based on their occurrence ratios in the benthic fauna by weighting them according to the concept of indicator groups (Simboura and Zenetos, 2002). AMBI, M-AMBI, and BENTIX indices are designed to evaluate the impact resulting from general stress factors and can not distinguish between natural and anthropogenic disturbances (Borja et al., 2003).

In order to detect and develop proper strategies for mitigating the effects of the discharge of anthropogenic chemical contaminants into marine ecosystems, we must assess and understand the ecological effects that affect habitat alteration and benthic marine communities. Hence, our study assessed the ecological quality of the Azini and Gwadar mangrove ecosystems in the Gulf of Oman using benthic quality indices including AMBI, M-AMBI, and BENTIX based on polychaete communities and determining the correlation of

these indices with environmental variables was investigated.

### Materials and methods

In the Iranian coastal zone, mangrove forests are distributed along 1250 km. Mangrove forests in Iran consist of two species of *Avicennia marina* and *Rhizophora mucronata*, that both species are found in the Azini estuary, but there is only *A. marina* in the Gwadar estuary. In the summer and winter of 2019, sediment samples were taken from mangrove forests in Gwadar and Azini estuaries (Fig. 1). Sediment samples were collected using a metal quadrat (25\*25\*25 cm) from 10 stations i.e. 5 stations for each estuary. The sediments were collected in three independent replicates for sediment variables and polychaetes. Polychaetes were sorted under a stereomicroscope and identified at the lowest taxonomic level possible. Measurements of total organic matter (TOM), total organic carbon (TOC), and the grain size composition of the sediment surface were performed using the ignition method (Heiri et al., 2001), the Walkley-Black (Walkley and Black, 1934) method,




Figure 2. AMBI ecological groups (EG) for the polychaete communities in mangrove ecosystems in the Gulf of Oman (the Gwadar and Azini estuaries).

Table 1. The threshold levels of three indices for benthic ecological quality status assessment.

Biotic index	Ecological quality status (EcoQs)				
	High	Good	Moderate	Poor	Bad
AMBI	0-1.2	1.2-3.3	3.3-4.3	4.3-5.5	5.5-7
M-AMBI	0.77-1	0.53-0.77	0.39-0.53	0.2-0.39	0.0-0.2
BENTIX	4.5-6	3.5-4.5	2.5-3.5	2-2.5	0.0-2

Note: The thresholds of AMBI and M-AMBI are referred to Borja et al. (2000), Borja and Muxika (2005), Muxika et al. (2007), Li et al. (2017), and the thresholds of BENTIX is referred to Simboura and Zenetos 92002)

and the hydrometer technique (Bouyoucos, 1962), respectively.

The AMBI and M-AMBI indices were used to assess the benthic ecological status of the study areas and calculated using the AMBI program (version 6.0; available at <http://ambi.azti.es>). The software divides polychaetes into five ecological groups (GI + GV) based on their sensitivity to environmental stress gradients. Following the newest AMBI species list (December 2020), most of the collected species were classified into different ecological groups (EGs).

The assignment of some species, including some native species, was based on expert opinion or on the assignment of other species in the same genus (Borja et al., 2008). To improve the reliability of the results, AMBI values with more than 20% unassigned individuals were removed from the AMBI analyses but included in the M-AMBI analyses (Borja and Muxika, 2005). M-AMBI values were calculated using Factor Analysis of AMBI, Shannon diversity ( $H'$ ), and species richness ( $S$ ) (Muxika et al., 2007). In order to assess ecological status, it was crucial to set an appropriate M-AMBI reference condition. Therefore, in the present study, the reference

conditions for the M-AMBI index proposed by Borja and Tunberg (2011) and Forchino et al. (2011), which increased the highest species diversity and richness values by 15% and decreased AMBI to half of the lowest value.

Bentix simplified the five ecological groups into two, the sensitive species group (GS), which includes GI and GII of the AMBI method, and the tolerant species group (GT), which includes GIII, GIV, and GV (Add-in v.1.0 version: (Simboura and Zenetos, 2002). Table 1 shows the different ecological quality statuses determined by AMBI, M-AMBI, and BENTIX.

To assess differences in benthic indices between regions, and seasons, we used a repeated measure PERMANOVA model (permutational multivariate analysis of variance), where 'region' and 'season' was fixed factors. Permutations of row data were unlimited and 9999 permutations were applied. Spearman correlation was done to examine the relationship between sediment parameters and benthic indices. The univariate and multivariate analyses were conducted using PRIMER Ver. 6 with the PERMANOVA add-on package and SPSS 21.0 software.

Table 2. List of polychaete species and their assigned ecological groups (EG) in mangrove ecosystems in the Gulf of Oman (the Gwadar and Azini estuaries).

Species	AMBI groups	BENTIX groups
<i>Aricidea fragilis</i>	I	I
<i>Armandia intermedia</i>	I	I
<i>Armandia</i> sp.	I	I
<i>Capitella aberranta</i>	V	II
<i>Ctenodrilus</i> sp.	NA	NA
<i>Glycinde</i> sp.	II	I
<i>Heteromastus</i> sp.1	IV	II
<i>Heteromastus</i> sp.2	IV	II
<i>Heteromastus</i> sp.3	IV	II
<i>Heteromastus</i> sp.4	IV	II
<i>Leonnates indicus</i>	III	II
<i>Lepidonotus purpureus</i>	II	I
<i>Lepidonotus</i> sp.	II	I
<i>Levinsenia gracilis</i>	III	II
<i>Linopherus hirsuta</i>	IV	II
<i>Marphysa sanguinea</i>	II	I
<i>Marphysa</i> sp.	II	I
<i>Mediomastus warrenae</i>	NA	NA
<i>Melinna monocerooides</i>	III	II
<i>Namalycastis</i> sp.	NA	NA
<i>Neanthes glandicincta</i>	I	I
<i>Paleaequor</i> sp.	I	I
<i>Perinereis horsti</i>	III	II
<i>Perinereis nuntia</i>	III	II
<i>Phyllodoce</i> sp.	II	I
<i>Prionospio</i> sp.	NA	NA
<i>Questa riseri</i>	II	I
<i>Scolecopsis</i> sp.	III	II
<i>Scolecopsis squamata</i>	III	II
<i>Sigambra sundarbanensis</i>	IV	II
<i>Simplisetia erythraeensis</i>	III	II
<i>Syllis gracilis</i>	II	I
<i>Tylonereis bogoyawlensky</i>	III	II

\*NA: Not assigned

## Results

The collected species were categorized into five ecological groups to calculate the AMBI and M-AMBI indices (Fig. 2, Table 2). The highest number of species was in the ecological group (I) in Azini (84.9% of the total species in Azini). The lowest number of species were ecological groups (V) in the Gwadar. During two sampling seasons, ecological groups (I), (III), and (IV) included the most dominant species in the study regions.

The lowest and highest values of the AMBI index in all three regions were  $0.8 \pm 0.09$  and  $3.2 \pm 0.5$ , respectively, at Azini and Gwadar in summer. According to AMBI, Gwadar, and Azini were of good to high status (Fig. 3), and the results showed a

significant difference between regions ( $P < 0.05$ ). The AMBI index revealed that summer has better ecological quality, but there was no significant difference between seasons in both areas ( $P > 0.05$ ) (Table 3). M-AMBI showed that the ecological status of the study regions was good status and it ranged from  $0.59 \pm 0.08$  to  $0.52 \pm 0.13$  (Fig. 4). No significant difference was found between the two regions and seasons ( $P > 0.05$ ) (Table 3).

Two sensitive and tolerant ecological groups were used to calculate the BENTIX index (Fig. 5). The most sensitive species were found in Azini during summer (GS, 86.6%). The most tolerant species were observed in Gwadar during winter (GT, 67%).

Table 3. The results of PERMANOVA for comparing AMBI, M-AMBI and BENTIX across regions and seasons in mangrove ecosystems in the Gulf of Oman (the Gwadar and Azini estuaries). Factors: region (levels: Gwadar and Azini) and season (levels: winter and summer).

AMBI	df	Ms	Pseudo-F	P
Region	2	8638.4	2.0915	0.048
Season	1	6560.5	1.8017	0.259
Region × Season	1	5128.5	1.1307	0.438
Residual	62	2304.8		
Total	71			
M-AMBI	df	Ms	Pseudo-F	P
Region	2	1055.4	0.81601	0.567
Season	1	309.17	0.41735	0.142
Region × Season	1	150.01	0.2025	0.795
Residual	62	803		
Total	71			
BENTIX	df	Ms	Pseudo-F	P
Region	2	2825.3	1.3381	0.037
Season	1	787.02	0.60774	0.561
Region × Season	1	834.85	0.64468	0.519
Residual	62	582.58		
Total	71			




Figure 3. The AMBI values of polychaete communities in mangrove ecosystems in the Gulf of Oman (the Gwadar and Azini estuaries).




Figure 4. The M-AMBI values of polychaete communities in mangrove ecosystems in the Gulf of Oman (the Gwadar and Azini estuaries).

Table 4. Variations in environmental data (mean  $\pm$  SD) between regions and seasons in mangrove ecosystems in the Gulf of Oman (the Gwadar and Azini estuaries). (TOM = Total Organic Matter; TOC = Total Organic Carbon).

Gwadar	Winter	Summer
TOM	0.51 $\pm$ 0.09	0.71 $\pm$ 0.06
TOC	0.91 $\pm$ 0.3	1.23 $\pm$ 0.4
Silt/Clay (%)	1.13 $\pm$ 0.54	1.93 $\pm$ 1.5
Azini	Winter	Summer
TOM	0.78 $\pm$ 0.1	0.57 $\pm$ 0.08
TOC	0.99 $\pm$ 0.2	0.84 $\pm$ 0.3
Silt/Clay (%)	1.96 $\pm$ 0.91	1.62 $\pm$ 0.88




Figure 5. BENTIX ecological groups (EG) for the polychaete communities in mangrove ecosystems in the Gulf of Oman (the Gwadar and Azini estuaries).




Figure 6. The BENTIX values of polychaete communities in mangrove ecosystems in the Gulf of Oman (the Gwadar and Azini estuaries).

BENTIX index values ranged from  $2.8\pm 0.32$  to  $5\pm 0.67$  and Gwadar and Azini had the lowest and the highest values in summer, respectively. Using the BENTIX index, two regions were moderate to high status (Fig. 6). There was a significant difference in

ecological quality between Gwadar and Azini ( $P < 0.05$ ). The mean values of BENTIX did not differ significantly between winter and summer ( $P > 0.05$ ) (Table 3). Table 4 shows the physicochemical characteristics of the sediment during winter and

Table 5. Spearman correlation between biotic indices and sediment parameters in mangrove ecosystems in the Gulf of Oman (the Gwadar and Azini estuaries). (\*,  $P < 0.05$ ; \*\*,  $P < 0.01$ ).

	AMBI	M-AMBI	BENTIX	Silt/Clay	TOC
AMBI					
M-AMBI	-0.59				
BENTIX	-0.97	0.66			
Silt/Clay	-0.26	<b>-0.51*</b>	0.07		
TOC	<b>0.61*</b>	<b>-0.95**</b>	<b>-0.73*</b>	0.58	
TOM	0.46	<b>-0.91**</b>	-0.46	0.36	0.75

summer from the Gwadar and Azini estuaries. The values of M-AMBI showed the best response to increasing concentrations of TOM ( $r = -0.91$ ,  $P < 0.01$ ) and TOC ( $r = -0.95$ ,  $P < 0.01$ ). The AMBI and BENTIX values showed a significant correlation with TOC ( $r = -0.61$ ,  $P < 0.05$ , and  $r = -0.73$ ,  $P < 0.05$  respectively). Conversely, AMBI did not show significant correlations with TOM, and Silt/Clay (Table 5).

## Discussion

AMBI, M-AMBI, and BENTIX were used to assess the ecological quality of Gwadar and Azini mangroves in the Gulf of Oman. There were noteworthy differences between the results of the three biological indices. In comparison to AMBI and M-AMBI, the BENTIX showed a much lower ecological status. The indices showed different EcoQ classifications in the two regions, including "high" to "moderate" conditions in BENTIX and "high" to "good" conditions in AMBI, and "good" conditions in M-AMBI. Some studies report inconsistent classifications of EcoQs based on different benthic indices for the same study area (Simboura et al., 2014; Pelletier et al., 2018; Maghsoudlou et al., 2020; Yan et al., 2020; Xu et al., 2021; Dong et al., 2021). These differences could be due to (1) these indicators are designed to evaluate European coastal waters' ecosystems (Qiu et al., 2018; Xu et al., 2021), (2) there are different boundary limits (Simboura and Reizopoulou, 2008; Borja and Tunberg, 2011; Gamito et al., 2012; Sun et al., 2018; Ni et al., 2019), (3) in the various ecological groups, each index input is weighted differently (Simboura and Reizopoulou, 2008; Sun et al., 2018), and (4) worldwide,

macrobenthos is composed differently in marine ecosystems, and they may respond differently to disturbances (Borja et al., 2000; Pelletier et al., 2018; Mulik et al., 2020; Dong et al., 2021). The Benthic classification method yields a wider high-status range (4.5-6) than the AMBI (0-1.2), which often classifies good-status sites as high-status.

In the current study, the TOC content of sediment was significantly correlated with AMBI, M-AMBI, and BENTIX indices. According to the results, the BENTIX index showed a significant negative correlation with the TOC content of the sediment. Although organic matter provides food for benthic fauna in surface sediments, excessive amounts of organic enrichment can lead to oxygen depletion as well as toxic by-products, resulting in reduced species richness, abundance, and biomass of benthic fauna that are closely associated with bottom sediments (Hyland et al., 2005). The TOC content of sediment was higher in Gwadar mangroves compared to Azini mangroves. The BENTIX index was classified the Gwadar mangroves in "moderate" status, while the Azini mangroves classified as "good" to "high" status. Therefore, the TOC content of sediment can be an important descriptor and proxy for evaluating the benthic status of mangrove ecosystems in relation to ecological indices. The M-AMBI index showed the highest correlation with TOM and TOC content of sediments. Therefore, according to the results, the AMBI, M-AMBI, and BENTIX indices can be considered suitable indicators for enriching organic matter in sediments. Some studies observed correlations between organic matter and AMBI, M-AMBI and BENTIX indices (Borja et al., 2008;

Caglar and Albayrak 2012; Umehara et al. 2019; Medeiros et al. 2021; Munari et al. 2022; Sarathy et al. 2022). However, according to Hu et al. (2018), benthic indices are not correlated with organic matter, which may explain its overestimation of benthic quality.

The Azini mangroves were dominated by sensitive species (EG I), while the Gwadar mangroves were dominated by tolerant species (EG III) and second opportunistic species (IV). Apart from the absence of industries in the coastline region, the presence of two types of trees, including *A. marina* and *R. mucronata*, may have contributed to the "high" ecological status in Azini mangroves. According to Delfan et al. (2021), the low species richness of the macrofauna may be explained by a low diversity of mangrove trees. *Avicennia marina* is the only mangrove species in Gwadar. In addition, the wastewater from shrimp farming centers may have disturbed the Gwadar mangroves.

In the current study, because of a lack of knowledge, some of the collected species, especially some native species, could not be assigned to the AMBI species list, resulting in partly affecting index evaluation results. Lastly, it is important not to ignore the characteristics of the local ecosystem when assessing its status of the local ecosystem. Hence, no one biotic index is the most suitable, and when doing ecological assessments, it is more reliable to combine the results from several indices.

## Conclusion

The indices showed different EcoQ classifications in the two regions studied, including "high" to "moderate" status in BENTIX and "high" to "good" status in AMBI, and "good" status in M-AMBI. Sensitive species were dominant in the Azini mangroves, while resistant species were dominant in the Gwadar mangroves. There were significant correlations between benthic indices and the TOC content of sediments. Therefore, TOC can be a suitable descriptor and indicator to evaluate the quality of mangrove ecosystems in relation to

benthic indices. Also, the highest correlation was observed between M-AMBI and sediment parameters. Our results showed that several benthic indices are needed for assessing the benthic quality of ecosystems; as well as suggest useful insights on identifying the key drivers of polychaete communities and benthic indices in mangrove ecosystems.

## Acknowledgments

We gratefully acknowledge the financial support provided by Department of Marine Biology, Faculty of Natural Resources and Marine Sciences, Tarbiat Modares University, Noor, Iran.

## References

- Alongi D.M. (2002). Present state and future of the world's mangrove forests. *Environment Conservation*, 29(3): 331-349.
- Alvarez-Aguilar A., Rodríguez-Villanueva V., Macías-Zamora J.V., Ramirez-Alvarez N., Hernandez-Guzman F.A. (2017). Spatio-temporal analysis of benthic polychaete community structure in the north-western coast of Baja California, Mexico. *Journal of the Marine Biological Association of the United Kingdom* 97(5): 993-1005.
- Borja A., Dauer D.M., Díaz R., Llanso R.J., Muxika I., Rodriguez J.G., Schaffner L. (2008). Assessing estuarine benthic quality conditions in Chesapeake Bay: A comparison of three indices. *Ecological Indicators*, 8: 395-403.
- Borja A., Franco J., Pérez V. (2000). A marine biotic index to establish the ecological quality of soft-bottom benthos within European estuarine and coastal environments. *Marine Pollution Bulletin*, 40: 1100-1114.
- Borja Á., Marín S., Núñez R., Muxika I. (2014). Is there a significant relationship between the benthic status of an area, determined by two broadly-used indices, and best professional judgment? *Ecological Indicators*, 45: 308-312.
- Borja A., Muxika I. (2005). Guidelines for the use of AMBI (AZTI's Marine Biotic Index) in the assessment of the benthic ecological quality. *Marine Pollution Bulletin*, 50: 787-789.
- Borja A., Muxika I., Franco J. (2003). The application of

- a Marine Biotic Index to different impact sources affecting soft-bottom benthic communities along European coasts. *Marine Pollution Bulletin*, 46:835-845.
- Borja A., Tunberg B.G. (2011). Assessing benthic health in stressed subtropical estuaries, eastern Florida, USA using AMBI and M-AMBI. *Ecological Indicators*, 11: 295-303.
- Bouyoucos G.J. (1962). Hydrometer Method Improved for Making Particle Size Analyses of Soils. *Agronomy Journal*, 54(5): 464-465.
- Caglar S., Albayrak S. (2012). Assessment of ecological quality status of Küçükçekmece Bay (Marmara Sea) by applying BENTIX, AMBI, BOPA and BO2A biotic indexes. *Mediterranean Marine Science*, 13(2): 198-207.
- Dauvin J.C., Andrade H., de-la-Ossa-Carretero J.A., Del-Pilar-Ruso Y., Riera R. (2016). Polychaete/amphipod ratios: An approach to validating simple benthic indicators. *Ecological Indicators*, 63: 89-99.
- Delfan N., Shojaei M.G., Naderloo R. (2021). Patterns of structural and functional diversity of macrofaunal communities in a subtropical mangrove ecosystem. *Estuarine, Coastal and Shelf Science*, 252: 107288.
- Dong J-Y., Sun X., Zhang Y., Zhan Q., Zhang X. (2021). Assessing benthic habitat ecological quality using four benthic indices in the coastal waters of Sanshandao, Laizhou Bay, China. *Ecological Indicators*, 129: 107980.
- Duke N., Meynecke J-O., Dittmann S., Ellison A., Anger K. (2007). A world without mangroves? *Science (New York, NY)*, 317(5834): 41-42.
- Field C., Osborn J., Hoffmann L., Polsenberg J., Ackerly D. (1998). Mangrove biodiversity and ecosystem function. *Global Ecology and Biogeography Letters*, 7(1): 3-14.
- Forchino A., Borja A., Brambilla F., Rodrigues J.G., Muxika I., Terova G., Saroglia M. (2011). Evaluating the influence of off-shore cage aquaculture on the benthic ecosystem in Alghero Bay (Sardinia, Italy) using AMBI and M-AMBI. *Ecological Indicators*, 11(5): 1112-1122.
- Gamito S., Patrício J., Neto J.M., Teixeira H., Marques, J.C. (2012). Feeding diversity index as complementary information in the assessment of ecological quality status. *Ecological Indicators*, 19: 73-78.
- Heiri O., Lotter A.F., Lemcke G. (2001). Loss on ignition as a method for estimating organic and carbonate content in sediments: reproducibility and comparability of results. *Journal of Paleolimnology*, 25: 101-110.
- Hu C., Shui B., Li W., Yang X., Zhang X. (2018). Assessing the ecological quality status of transplanted mangrove wetland in the Oujiang estuary, China. *Marine Pollution Bulletin*, 133: 1-8.
- Hyland J., Balthis L., Karakassis I., Magni P., Petrov, A. (2005). Organic carbon of sediments as an indicator of stress in the marine benthos. *Marine Ecology Progress Series*, 295: 91-103.
- Li B., Li X., Bouma T.J., Soissons L.M., Cozzoli, F. (2017). Analysis of macrobenthic assemblages and ecological health of Yellow River Delta, China, using AMBI & M-AMBI assessment method. *Marine Pollution Bulletin*, 119: 23-32.
- Maghsoudlou A., Momtazi F., Hashtroudi M.S. (2020). Ecological Quality Status (EcoQs) of Chabahar subtropical bay based on multimetric macrobenthos-indexes approach: Response of bio-indexes to sediment structural/pollutant variables. *Regional Studies in Marine Science*, 40: 101524.
- Medeiros C.R., Paiva F.F., Ligeiro R., Molozzi J., Melo A.S. (2021). Saline gradient drives functional nestedness of polychaete communities in tropical estuaries. *Estuarine, Coastal and Shelf Science*, 251: 107185.
- Mulik J., Sukumaran S., Srinivas T. (2020) Factors structuring spatio-temporal dynamics of macrobenthic communities of three differently modified tropical estuaries. *Marine Pollution Bulletin*, 150: 110767.
- Munari C., Borja A., Corinaldesi C., Rastelli E., Lo Martire M., Pitacco V., Mistri M. (2022). First Application of the AMBI Index to the Macrobenthic Soft-Bottom Community of Terra Nova Bay (Ross Sea, Southern Ocean). *Water*, 14(19): 2994.
- Muxika I., Borja Á., Bald J. (2007). Using historical data, expert judgement and multivariate analysis in assessing reference conditions and benthic ecological status, according to the European Water Framework Directive. *Marine Pollution Bulletin*, 55: 16-29.
- Ni D., Zhang Z., Liu X. (2019). Benthic ecological quality assessment of the Bohai Sea, China using marine biotic indices. *Marine Pollution Bulletin*, 142: 457-464.
- Pelletier P., Gillett D., Hamilton A., Grayson T., Hansen V., Leppo E.W., Weisberg S.B., Borja A. (2018).

- Adaptation and application of multivariate AMBI (M-AMBI) in US coastal waters.
- Qiu B., Zhong X., Liu X. (2018). Assessment of the benthic ecological status in the adjacent waters of Yangtze River Estuary using marine biotic indices. *Marine Pollution Bulletin*, 137: 104-112.
- Quarto A. (2005). Mangrove restoration-natural protection from natural disasters. *Biodiversity*, 6(1): 3-12.
- Sarathy P.P., Bharathidasan V., Murugesan P., Selvaraj P., Punniamorthy R. (2022). Seasonal carbonate system vis-à-vis pH and Salinity in selected tropical estuaries: Implications on polychaete diversity and composition towards predicting ecological health. *Oceanologia*, 64(2): 346-362.
- Schüller M., Ebbe B., Wägele J-W. (2009). Community structure and diversity of polychaetes (Annelida) in the deep Weddell Sea (Southern Ocean) and adjacent basins. *Marine Biodiversity*, 39: 95-108.
- Simboura N., Reizopoulou S. (2008). An intercalibration of classification metrics of benthic macroinvertebrates in coastal and transitional ecosystems of the Eastern Mediterranean ecoregion (Greece). *Marine Pollution Bulletin*, 56: 116-126.
- Simboura N., Tsapakis M., Pavlidou A., Assimakopoulou G., Pagou K. (2014). Assessment of the environmental status in the Hellenic coastal waters (Eastern Mediterranean): From the water framework directive to the marine strategy framework directive. *Mediterranean Marine Science*, 16: 46-64.
- Simboura N., Zenetos A. (2002). Benthic indicators to use in Ecological Quality classification of Mediterranean soft bottom marine ecosystems, including a new Biotic Index. *Mediterranean Marine Science*, 3(2): 77-111.
- Sun Y., Chen B., Wu H., Huang H., Ma Z., Tang K. (2018). Assessing benthic ecological status in subtropical islands, China using AMBI and Bentix indices. *Estuarine, Coastal and Shelf Science*, 207: 345-350.
- Tyler C., Kowalewski M. (2018). Regional surveys of macrobenthic shelf invertebrate communities in Onslow Bay, North Carolina, U.S.A. *Scientific Data*, 5(1): 1-7.
- Umehara A., Nakai S., Okuda T., Ohno M., Nishijima W. (2019). Benthic quality assessment using M-AMBI in the Seto Inland Sea, Japan. *Marine Environmental Research*, 148: 67-74.
- Walkley A., Black I.A. (1934). AN examination of the Degtjareff method for determining soil organic matter and a proposed modification of the chromic acid titration method. *Soil Science*, 37(1): 29-38.
- Xu J., Lu X., Liu X. (2021). Patterns of species and functional diversity of macrofaunal assemblages and the bioassessment of benthic ecological quality status in the southern Yellow Sea. *Marine Pollution Bulletin*, 171: 112784.
- Yan J., Sui J., Xu Y., Li X., Wang, H., Zhang B. (2020). Assessment of the benthic ecological status in adjacent areas of the Yangtze River Estuary, China, using AMBI, M-AMBI and BOPA biotic indices. *Marine Pollution Bulletin*, 153: 111020.