

Original Article

Protective effect of *Ocimum gratissimum* leaf aqueous extract on antioxidant capacity, immunocompetence, and disease resistance of Nile tilapia (*Oreochromis niloticus*) fingerlings against *Vibrio parahaemolyticus* infection

Rasaq Ibrahim^{1,2*}, Jehu Auta¹, John Ameh Adakole¹, Abdullateef Yusuf¹, Esther Yogbabaliat Yashim³, Abdullahi Bala Alhassan¹, Rilwanu Umar⁴, Harrison Charo-Karisa⁵, Femi John Fawole⁶, Hien Van Doan^{7,8}

¹Department of Biology, Ahmadu Bello University, Zaria, Nigeria.

²Department of Forestry and Fisheries, Kebbi State University of Science and Technology, Aliero, Nigeria.

³Department of Zoology, Ahmadu Bello University, Zaria, Nigeria.

⁴National Agricultural Extension and Research Liaison Services, Ahmadu Bello University, Zaria, Nigeria.

⁵WorldFish, Abbassa, Abou-Hammad, Sharkia, Egypt.

⁶Department of Aquaculture and Fisheries, University of Ilorin, Ilorin, Nigeria.

⁷Department of Animal and Aquatic Sciences, Faculty of Agriculture, Chiang Mai University, Chiang Mai 50200, Thailand.

⁸Functional Feed Innovation Centre (FuncFeed), Faculty of Agriculture, Chiang Mai University, Chiang Mai 50200, Thailand.

Abstract: Infectious diseases are major constraints on the aquaculture industry, and they frequently arise from stress factors associated with intensive aquaculture practices. This study was carried out to assess the protective effect of *Ocimum gratissimum* leaf aqueous extract on immunological responses, antioxidant capacity, and disease resistance of *Oreochromis niloticus* fingerlings against *Vibrio parahaemolyticus* infection. Five iso-nitrogenous (30.02%) diets containing 0 (control), 2, 4g, 8, and 16 g *Ocimum*-leaf extract (g/kg) were fed to triplicate groups of fish (initial weight 3.31±0.05g) for 90 days. After a 90-day feeding trial, test fish were injected with pathogenic *V. parahaemolyticus*, and survival was recorded over the next 14 days. The white blood cell (12.29±0.35×10³/μl), red blood cells (3.60±0.21×10³/μl), packed cell volume (27.76±0.08 g/dl), and haemoglobin (7.21±0.47 g/dl) were all highest in fish fed 8 g/kg of *Ocimum* leaf extract. The biochemical parameters, such as alkaline phosphatase (ALP), total protein, albumin, and globulin contents, increased significantly in *Ocimum* leaf extracts fed groups compared with the control; these groups also registered the lowest values of aspartate aminotransferase (AST), alanine aminotransferase (ALT), and malondialdehyde (MDA). Following a challenge with *V. parahaemolyticus*, the study indicated that *Ocimum* leaf extract reduced metabolic stress in infected fish compared with the control. There is a significant increase in glucose (101.00±2.89 mg/dl) levels and liver enzymes ALT (74.14±2.31 UI/L) and AST (49.00±1.73 UI/L) in the control group (0 g/kg), whereas the opposite was true for diets supplemented with *Ocimum* leaf extract. The *Ocimum* leaf extract group significantly enhanced immune parameters, including lysozyme, immunoglobulin (IgM), and Interleukin-1β (IL-1β), in the spleen, skin, and kidney. After infection, the supplemented diets containing *Ocimum* leaf extract showed significantly higher lysozyme and IgM activities in the spleen, skin, and kidney. The result of the challenge test with *V. parahaemolyticus* revealed that the highest survival (80%) was recorded in the group fed *Ocimum*-supplemented diets at 8 and 16 g/kg, and the lowest skin lesion morbidity (15%) was observed in fish fed 8 g/kg. This study revealed that the inclusion of *O. gratissimum* leaf extract in the diet of *O. niloticus* enhanced biochemical parameters and immune response, and improved resistance to *V. parahaemolyticus*.

Article history:

Received 3 June 2026

Accepted 25 June 2026

Available online 25 October 2026

Keywords:

Innate immunity

Cytokine

Nile tilapia

Pathogenic organism

Introduction

Aquaculture is currently one of the fastest-growing food-producing sectors worldwide and plays a pivotal role in global food security, human nutrition, and socioeconomic development (Abeyasinghe et al., 2025). The sector provides high-quality animal

protein while also generating employment opportunities and income for millions of people, particularly in developing countries (Liboureau et al., 2026; Nordhagen et al., 2026). Among cultured aquatic species, Nile tilapia (*Oreochromis niloticus*) is recognized as one of the most economically important

*Correspondence: Rasaq Ibrahim
E-mail: ibrahim.rasaq@ksusta.edu.ng

freshwater fish species worldwide, owing to its rapid growth rate, high consumer acceptance, adaptability to diverse environmental conditions, and efficient utilization of relatively inexpensive feeds (El-Sayed and Fitzsimmons, 2023; Vilhena Batista et al., 2026). Consequently, Nile tilapia contributes substantially to global aquaculture production and serves as an affordable and accessible source of animal protein in many tropical and subtropical regions (Koley et al., 2025). However, as with other intensively farmed fish species, the continuous expansion and intensification of tilapia farming have significantly increased fish vulnerability to infectious diseases, particularly under stressful rearing conditions and inadequate biosecurity practices (Chupani et al., 2026; Emam et al., 2025).

Disease outbreaks in tilapia culture systems frequently result in physiological stress, suppressed growth performance, reduced survival, and considerable economic losses for producers, while disease infestation and growth retardation remain among the principal factors limiting tilapia productivity in Nigeria and several other developing countries (Paredes-Trujillo and Mendoza-Carranza, 2022; Maezono et al., 2025; Ndaro et al., 2026). Among the bacterial diseases affecting cultured fish, vibriosis caused by *Vibrio* species is regarded as one of the most devastating infectious diseases in aquaculture worldwide (Sanches-Fernandes et al., 2022; Manchanayake et al., 2023; Camberos-Solano and Soto-Rodriguez, 2026). *Vibrio parahaemolyticus*, a Gram-negative opportunistic pathogen, can infect marine, brackish, and freshwater fish species, leading to septicemia, skin hemorrhage, tissue necrosis, and high mortality rates (Petrusic et al., 2025; Li et al., 2026). The disease is particularly severe in fry and fingerlings due to their relatively immature immune systems and lower resistance to pathogenic infections. Furthermore, *V. parahaemolyticus* is recognized as an important foodborne pathogen that poses substantial risks to both aquaculture production and public health (Zaher et al., 2021; ELTarabili et al., 2026). Therefore, improving the immune competence and disease resistance of cultured fish through nutritional

interventions and functional feed-based strategies has become an important, sustainable approach to enhancing fish health and supporting the long-term sustainability of aquaculture production systems.

In recent years, plant-derived feed additives and phytogetic compounds have attracted considerable attention in aquaculture as promising alternatives to synthetic antibiotics and chemotherapeutic agents (Ivanova et al., 2024; Bai et al., 2025; Pudota et al., 2025). Medicinal plants and their derivatives are rich sources of biologically active compounds, including flavonoids, phenolics, alkaloids, terpenoids, tannins, and essential oils, which exhibit antimicrobial, antioxidant, anti-inflammatory, and immune-stimulatory properties (Chakroborty et al., 2025; Hossain et al., 2026). Owing to these functional characteristics, dietary supplementation with phytogetic additives has been widely reported to enhance growth performance, feed utilization efficiency, antioxidant defense systems, immune responses, gut health, and disease resistance in cultured fish species (Hu et al., 2025; Ibrahim et al., 2025; Adhikary et al., 2026; Soares et al., 2026; Xi et al., 2026). Moreover, phytogetic feed additives are generally biodegradable, environmentally friendly, and safe for long-term use, making them attractive candidates for sustainable, antibiotic-free aquaculture production systems (Wang et al., 2024; Hassan et al., 2025).

Among the medicinal plants investigated for aquaculture applications, *Ocimum gratissimum* L. has emerged as a particularly promising phytogetic additive because of its broad spectrum of biological activities (Bandeira et al., 2017). *Ocimum gratissimum*, commonly known as clove basil or scent leaf, is an aromatic medicinal plant widely distributed throughout tropical and subtropical regions, especially in Africa and Asia (Ugbogu et al., 2021). The plant has long been used in traditional medicine for its diverse pharmacological properties, including antimicrobial, antioxidant, anti-inflammatory, and immune-modulatory effects (Kamelnia et al., 2023; Sagar et al., 2025). Previous studies have demonstrated that

O. gratissimum contains several important bioactive compounds, such as eugenol, thymol, linalool, methyl cinnamate, camphor, flavonoids, and terpenoids, which are largely responsible for its biological activities (Leite dos Santos et al., 2025; Odeniyi et al., 2025; Akanbi et al., 2026; Sharma et al., 2026). Additionally, Egba et al. (2014) reported that extracts of *O. gratissimum* enhanced both primary and secondary antibody responses and stimulated immune functions. In aquaculture, dietary administration of *O. gratissimum* has been associated with improvements in hematological parameters, antioxidant capacity, immune responses, and disease resistance (Abdel-Tawwab et al., 2018; Ballester et al., 2025). Collectively, these findings suggest that *O. gratissimum* could serve as an effective natural immunostimulant and antioxidant feed additive capable of enhancing fish health and resistance against pathogenic infections. However, despite the increasing interest in phytogenic feed additives, information regarding the protective effects of *O. gratissimum* leaf aqueous extract against *V. parahaemolyticus* infection in Nile tilapia remains limited, particularly concerning its influence on hematological responses, oxidative stress biomarkers, innate immune parameters, and post-challenge disease resistance. Therefore, the present study was conducted to evaluate the protective effects of dietary *O. gratissimum* leaf aqueous extract on hematological responses, serum biochemical parameters, antioxidant status, innate immune responses, and disease resistance of Nile tilapia fingerlings challenged with *V. parahaemolyticus*.

Materials and Methods

Collection of leaf samples: Fresh leaves of *O. gratissimum* were collected from Saye Village, located in Zaria Local Government Area, Kaduna State, Nigeria. The plant material was taxonomically authenticated at the Herbarium Unit, Department of Botany, Faculty of Life Sciences, Ahmadu Bello University, where a voucher specimen was deposited under the accession number ABU01285.

Preparation of aqueous extract of *O. gratissimum*:

The collected leaves of *O. gratissimum* were dried and ground into a fine powder. Briefly, 2500 g of the powdered sample was macerated in 2.5 L of distilled water at room temperature for 24 h. The resulting mixture was subsequently filtered through Whatman No. 1 filter paper to remove particulate materials. The filtrate was then concentrated in a water bath maintained at 40°C until a semi-solid extract was obtained. The prepared aqueous extract was transferred into sterile sample bottles and stored under appropriate conditions until further use, following the method described by Salau et al. (2013).

Determination of phytochemical components of

***O. gratissimum* leaf aqueous extract:** The phytochemical composition of *O. gratissimum* leaf aqueous extract was determined using Gas Chromatography–Mass Spectrometry (GC-MS) analysis conducted at the Multiuser Science Laboratory, Department of Chemistry, Ahmadu Bello University. The analysis was performed to identify and characterize the bioactive compounds present in the aqueous extract (Table 1), while the corresponding total ion chromatogram is presented in Figure 1.

GC-MS analysis was carried out using a 7890B GC System coupled with a 5977A Mass Selective Detector (Agilent Technologies, USA). Prior to analysis, the semi-solid extract of *O. gratissimum* was diluted in methanol, and 2 µL of the prepared solution was injected into the GC-MS system using a microsyringe. The operating conditions of the GC-MS system were as follows: initial oven temperature of 70°C with an equilibration time of 1 min, followed by a programmed increase at 5°C min⁻¹ to 250°C, then maintained for 1 min, and finally increased at 30°C min⁻¹ to 300°C. The maximum oven temperature was set at 325°C, with a post-run temperature of 50°C. Helium was used as the carrier gas under split injection mode. The injector temperature was maintained at 250°C with a pressure of 11.089 psi, a total flow rate of 19.204 mL min⁻¹, a septum purge flow of 3 mL min⁻¹, and a purge flow to the split vent of 15 mL min⁻¹ at 0.75 min. The hold-up time and

Retention Time (minute)	% Peak Area	Name of Compounds	Reference
5.80	0.17	Eucalyptol	27465
6.03	0.06	Oxirane, tetradecyl-	102576
6.39	0.21	Benzaldehyde, 2-hydroxy-	10083
8.00	0.07	Hexanoic acid, methyl ester	13783
10.89	0.29	Methenamine	18654
12.26	0.28	3-Hexyne-2,5-diol, 2,5-dimethyl-	20272
12.89	0.85	Thymol	24339
17.75	0.36	Cyclohexane,1-ethenyl-1-methyl-2,4-bis(1-methylethenyl)-	68686
18.53	0.30	Methyl 8-methyl-decanoate	65001
20.17	0.15	Fluoroacetic acid, dodecyl ester	107808
20.35	0.07	2-Methyltetracosane	207500
23.37	0.22	Tridecanoic acid, methyl ester	91455
24.88	0.30	Heptadecyl trifluoroacetate	207045
25.02	0.11	Hexadecane, 1-chloro-	121205
25.47	0.19	Phenol, 2,6-diamino-	10528
25.70	0.23	3-Pentanone, 2,4-dimethyl-, O-methyloxime	20882
26.71	0.13	Oxirane, decyl-	51291
27.81	2.88	Pentadecanoic acid, 14-methyl-, methyl ester	130841
28.69	0.29	1,2-Benzenedicarboxylic acid, butyl 2-methylpropyl ester	138186
29.17	0.23	Pentafluoropropionic acid, tetradecyl ester	213156
31.06	0.98	1-Hexadecanol	104431
31.24	1.25	9,12-Octadecadienoic acid (Z,Z)-, methyl ester	153891
31.36	5.43	9-Octadecenoic acid (Z)-, methyl ester	155751
31.64	0.55	Phytol	155851
31.85	2.76	Methyl stearate	157880
32.64	0.09	7,11-Hexadecadienal	98680
32.81	0.28	1-Isopropoxy-2,2, 3-trimethylaziridine (anti)	20887
33.08	0.36	17-Pentatriacontene	265113
33.58	0.06	Pentanoic acid, 10-undecenyl ester	115375
34.02	0.07	2-Methyl-Z,Z-3,13-octadecadienol	140253
35.56	0.34	Undecanoic acid, 2-methyl-	65005
35.68	0.16	9,12-Octadecadienoyl chloride, (Z,Z)-	157778
35.81	0.17	Oleic Acid	142069
36.13	0.44	2-Methyl-Z,Z-3,13-octadecadienol	140253
36.71	0.10	Cyclohexanemethanol	7618
36.83	0.62	Hexanedioic acid, bis(2-ethylhexyl) ester	220809
37.81	60.57	9,17-Octadecadienal, (Z)-	125003
38.36	18.41	Benzoic acid, 4-amino-2-propoxy-, 2-(diethylamino)ethyl ester	153490

Table 1. GC-MS profile of *Ocimum gratissimum* leaf aqueous extract.

carrier gas flow rate were 1.2386 min and 1.0 mL min⁻¹, respectively.

Experimental diet preparation: The feed ingredients used in diet formulation were purchased from Sabon-Gari Market in Zaria, Kaduna State, Nigeria. All ingredients were cleaned, processed, and finely milled prior to diet preparation. Five experimental diets (Diets 1-5) were formulated to be isonitrogenous, containing 29.78-30.39% crude protein (Table 2), and supplemented with different inclusion levels of *O. gratissimum* leaf aqueous extract at 0, 2, 4, 8, and 16 g kg⁻¹ diet.

The basal diet (0 g kg⁻¹ *O. gratissimum* extract; control diet) was prepared using yellow maize, wheat bran, soybean meal, fish meal, corn starch, chicken intestine, palm oil, vitamin–mineral premix, lysine, and methionine. The ingredients were thoroughly mixed with water to form a homogeneous dough, then pelletized, air-dried, and subsequently crushed into pellets of appropriate size for the experimental fish. The supplemented diets were prepared using the same basal formulation with the respective inclusion levels of *O. gratissimum* leaf aqueous extract.

During the feeding trial, fish were fed the

Table 2. Composition of the experimental diets with levels of *Ocimum gratissimum* leaf extract.

Ingredients (kg)	<i>Ocimum</i> leaf extract levels (g/kg diet)				
	0	2	4	8	16
Yellow corn	350	350	350	350	350
Soya bean meal	200	200	200	200	200
Wheat bran	60	60	60	60	60
Corn starch	100	100	100	100	100
Fish meal	75.0	75.0	75.0	75.0	75.0
Chicken intestine	150	150	150	150	150
Palm oil	30	30	30	30	30
Premix ^a	15.0	15.0	15.0	15.0	15.0
Lysine	10	10	10	10	10
Methionine	10	10	10	10	10
Proximate composition (%)					
Crude protein	29.78	30.07	29.84	30.00	30.39
Crude lipid	8.23	8.12	8.50	8.64	8.23
Nitrogen free extract	34.78	33.22	33.71	34.02	34.56
Crude fibre	11.06	12.06	12.00	12.00	12.67
Ash	6.99	7.01	6.98	7.79	7.18

Premix^a: vitamin and mineral premix (IU or mg/kg). Vitamin A, 4800 IU; vitamin D3, 2400 IU; vitamin E, 4000 mg; vitamin K: 800 mg; Vitamin B1: 400mg; vitamin B2, 1600 mg; vitamin B6, 600 mg; vitamin B12, 4 mg; pantothenic acid, 4000 mg; nicotinic acid, 8000mg; folic acid, 400 mg; biotin, 20 mg, Manganese, 22000 mg; zinc, 22000 mg; iron, 12000 mg; copper, 4000 mg; iodine, 400 mg; selenium, 400mg; cobalt, 4.8 mg.




Figure 1. Total ion chromatogram of *Ocimum gratissimum* leaf aqueous extract

experimental diets to apparent satiation three times daily for 90 days. The experimental diets were weighed before the commencement and at the end of the feeding trial to estimate feed intake and determine the amount of feed administered to each experimental tank.

Experimental fish and rearing conditions: A total

of 800 healthy fingerlings of Nile tilapia were obtained from a commercial fish farm in Kano State, Nigeria. The fish were transported in 50-L plastic containers to the Fisheries Research Laboratory, Department of Biology, Faculty of Life Sciences, Ahmadu Bello University. Upon arrival, the fish were acclimatized in holding tanks containing dechlorinated tap water and fed a commercial diet (Coppens) at 5% of body weight for two weeks prior to the commencement of the feeding trial.

After acclimation, fish with an initial mean body weight ranging from 3.28±0.03 to 3.36±0.05 g were randomly distributed into plastic experimental tanks (30.0×20.0×25.0 cm) containing 35 L of water. Each tank was stocked with 15 fish, and each dietary treatment was conducted in triplicate, resulting in 45 fish per treatment group. The fish were assigned to one of five experimental diets containing 0 (control), 2, 4, 8, or 16 g kg⁻¹ of *O. gratissimum* leaf aqueous extract.

Before the initiation of the feeding trial, all fish were batch-weighted and measured to determine their initial body weight and total length using a top-loading digital balance (AT. M A 123). Final measurements were similarly recorded at the end of the 90-day experimental period.

Determination of hematological parameters: At the end of the 90-day feeding trial, three fish were randomly sampled from each experimental tank for hematological analysis. Prior to sampling, fish were fasted for 24 h to minimize postprandial variations in blood biochemical and hematological parameters. Blood samples were collected from the heart and caudal peduncle using 1-mL hypodermic syringes pre-rinsed with ethylenediaminetetraacetic acid (EDTA) as an anticoagulant.

For serum collection, nine fish per treatment were euthanized using buffered benzocaine solution (>250 mg L⁻¹). Fish were maintained in the anesthetic solution for approximately 10 min following cessation of opercular movement to ensure complete euthanasia. Hematological analyses were subsequently performed using a Biobase BK6100 Hematology Analyzer (Biobase, China) operated in whole-blood and predilute modes.

The hematological parameters evaluated included white blood cell count (WBC), red blood cell count (RBC), hemoglobin concentration (HGB), and packed cell volume (PCV). In addition, mean corpuscular volume (MCV), mean corpuscular hemoglobin (MCH), and mean corpuscular hemoglobin concentration (MCHC) were calculated according to the methods described by Dacie and Lewis (1975) using the formula of $MCV (\mu m^3) = Ht / (RBC's) \times 10$, and $MCH (pg) = Hb / (RBC's) \times 10$.

Serum biochemical assessment for liver enzymes:

Serum biochemical parameters, including alanine aminotransferase (ALT), aspartate aminotransferase (AST), alkaline phosphatase (ALP), total protein, albumin, glucose, and globulin, were determined using commercial colorimetric assay kits (Randox Laboratories Ltd., London, UK) following the manufacturer's instructions. The absorbance values of the standards and samples were measured relative to a reagent blank using a spectrophotometer at 630 nm.

Sample preparation for oxidative biomarkers and immune parameters:

Skin mucus samples were collected by gently scraping the dorsal body surface of the experimental fish using sterile glass slides.

Following mucus collection, the fish were dissected along the left lateral side to expose the abdominal cavity, and tissue samples, including liver, spleen, and kidney, were carefully excised. The collected tissue samples and skin mucus were each homogenized in physiological saline solution and then centrifuged at $6000 \times g$ for 20 min at 4°C. The resulting supernatants were carefully collected and stored under appropriate conditions for subsequent analyses of oxidative stress biomarkers and immune-related parameters.

Determination of oxidative stress biomarkers:

Liver tissues of *O. niloticus* were collected and immediately perfused with chilled physiological saline solution (0.9% NaCl). The tissues were homogenized in physiological saline, filtered, and centrifuged at 1500 rpm for 20 min at 4°C. The resulting clear supernatants were collected and used for analyses of oxidative stress biomarkers. Total protein content in the supernatant was determined and expressed relative to the wet tissue weight. The oxidative stress biomarkers analyzed included reduced glutathione (GSH), superoxide dismutase (SOD), catalase (CAT), and malondialdehyde (MDA), following the methods described by Nishikimi et al. (1972), Beutler et al. (1963), Aebi (1984), and Ohkawa et al. (1979), respectively. Commercial assay reagents and kits were obtained from Biodiagnostic Co.

Reduced glutathione (GSH) levels were determined spectrophotometrically using a Photometer 5010 (BM Co., Germany) according to the method of Beutler et al. (1963). Catalase (CAT) activity was assessed by measuring the rate of hydrogen peroxide decomposition at 240 nm, following the procedure described by Aebi (1984). Superoxide dismutase (SOD) activity was determined by diluting 0.1 mL of tissue homogenate with 0.9 mL distilled water to obtain a 1:10 dilution. Subsequently, 0.2 mL of the diluted sample was mixed with 2.5 mL of 0.05 M carbonate buffer, followed by the addition of 0.3 mL of 0.3 mM adrenaline solution. The reference mixture consisted of 2.5 mL of carbonate buffer, 0.3 mL of adrenaline solution, and 0.2 mL of

distilled water. Absorbance was measured at 480 nm from 30 to 150 s, and SOD activity was calculated based on the percentage inhibition of adrenaline auto-oxidation.

Lipid peroxidation was estimated by measuring thiobarbituric acid reactive substances (TBARS) as described by Ohkawa et al. (1979). Briefly, 2 mL of 15% trichloroacetic acid and 2 mL of thiobarbituric acid solution were added to 100 μ L of tissue homogenate. The mixture was incubated in a water bath at 80°C for 30 min, cooled to room temperature, and centrifuged at 3000 rpm for 10 min. The absorbance of the resulting supernatant was measured at 535 nm using a spectrophotometer. The concentration of TBARS was expressed as nmol/mg protein.

Determination of immune parameters: Immune-related parameters were assessed in the spleen, skin, and kidney tissues of the experimental fish using enzyme-linked immunosorbent assay (ELISA) kits according to the manufacturers' instructions. The evaluated immune parameters included lysozyme activity (Fish LZM CK-LAB Kit), immunoglobulin M (IgM; Fish IgM CK-LAB Kit), and interleukin-1 β (IL-1 β ; Interleukin 1 Beta FineTest ELISA Kit). The lysozyme and IgM assay kits were obtained from Shanghai Coon Koon Biotech Co., Ltd. All assays were performed according to the manufacturers' protocols.

Challenge test of experimental fish with *V. parahaemolyticus*: At the end of the 90-day feeding trial, the experimental fish were challenged with a virulent strain of *V. parahaemolyticus*. The bacterial strain was obtained from the Department of Veterinary Microbiology, Ahmadu Bello University. The bacterium was cultured on blood agar plates and incubated at 37°C for 24 h prior to the challenge test. A preliminary experiment was conducted to determine the median lethal dose (LD₅₀) of *V. parahaemolyticus*. Fish were intraperitoneally injected with 0.1 mL of bacterial suspensions containing different concentrations of *V. parahaemolyticus* (1.5×10^8 , 3.5×10^8 , 6×10^8 , and 9×10^8 CFU mL⁻¹). Bacterial

density was quantified spectrophotometrically at an optical density (OD) of 450 nm. The results of the preliminary trial indicated that 0.1 mL containing 3.7×10^8 CFU mL⁻¹ represented the 96-h LD₅₀ dose.

For the challenge experiment, 10 healthy fish were randomly selected from each treatment tank and intraperitoneally injected with 0.1 mL phosphate-buffered saline (PBS) containing 3.7×10^8 CFU mL⁻¹ of *V. parahaemolyticus*. Prior to bacterial challenge, fish were fasted for 24 h, after which feeding with the respective experimental diets was resumed. Following infection, fish were monitored daily for 14 days to record mortality and observe abnormal clinical signs. At the end of the challenge period, the severity of skin hemorrhage and lesions was evaluated using a point-based grading system as described by Liu et al. (2018).

Water quality monitoring: Water quality parameters were monitored throughout the experimental period to ensure suitable rearing conditions for the fish. Water temperature (27.31-27.48°C), pH (7.29-7.37), electrical conductivity (206.67 μ S cm⁻¹), and total dissolved solids (103.33 - 105.33 ± 1.26 mg L⁻¹) were measured daily using a Hanna multiparameter instrument (Model No. 626 973-01). Dissolved oxygen concentration (7.13 ± 0.17 mg L⁻¹) was determined according to the standard methods described by APHA (2007).

Data analysis: All data were initially tested for normality and homogeneity of variances using the Shapiro-Wilk and Levene's tests, respectively. Subsequently, the collected data were analyzed using one-way analysis of variance (ANOVA) in R (version 4.1.2). When significant differences were detected, Tukey's multiple comparison test was applied to compare treatment means. Differences were considered statistically significant at $P < 0.05$. In addition, dependent t-tests were performed to compare parameters before and after the bacterial challenge. Graphs were generated using GraphPad Prism software (version 9.0).

Results

Hematological parameters: The hematological




Figure 2. Haematological parameters. (a) White blood cell, (b) Red blood cell, (c) Packed cell volume, (d) Haemoglobin, (e) Mean corpuscular volume, and (f) Mean corpuscular haemoglobin of *Oreochromis niloticus* fed a diet with different levels of *Ocimum gratissimum* leaf extract pre- and post-challenged with *Vibrio parahaemolyticus*. Values are mean $\pm$ SE, bars with different letters (a, b, c) differ significantly at $P < 0.05$. Upper case letters (A and B) indicated a significant difference at $P < 0.05$ between pre- and post-challenging with *V. parahaemolyticus*.

parameters of *O. niloticus* fed diets supplemented with different levels of *O. gratissimum* leaf extract before and after challenge with *V. parahaemolyticus* are presented in Figure 2. White blood cell (WBC) counts were significantly higher ($P < 0.05$) in all dietary treatment groups compared with the control group

under both pre- and post-challenge conditions, with the highest values observed in fish fed 8 g kg⁻¹ *Ocimum* leaf extract. Furthermore, WBC levels increased significantly following the bacterial challenge, indicating an enhanced immune response against *V. parahaemolyticus*. Similarly, red blood cell

(RBC) counts were significantly elevated ($P<0.05$) in fish fed *Ocimum*-supplemented diets compared with the control group prior to challenge. However, no significant differences ($P>0.05$) in RBC values were observed among the dietary groups following bacterial infection.

Hemoglobin (HGB) was also significantly higher ($P<0.05$) in fish receiving *Ocimum* leaf extract-supplemented diets than in the control group during the pre-challenge period. The highest HGB value (7.21 ± 0.47 g dL⁻¹) was recorded in fish fed 8 g kg⁻¹ *Ocimum* leaf extract, which was comparable to the values observed in fish fed 16 g kg⁻¹ (6.79 ± 0.17 g dL⁻¹), 4 g kg⁻¹ (6.65 ± 0.08 g dL⁻¹), and 2 g kg⁻¹ (5.89 ± 0.06 g dL⁻¹), but significantly higher than that of the control group (5.37 ± 0.47 g dL⁻¹). Following challenge with *V. parahaemolyticus*, a marked reduction in HGB concentration was observed in the control group, decreasing from 5.37 ± 0.47 to 3.43 ± 0.69 g dL⁻¹, whereas no significant differences ($P>0.05$) were detected among the treatment groups. Overall, WBC, RBC, PCV, and HGB values increased progressively with increasing dietary inclusion levels of *Ocimum* leaf extract up to 8 g kg⁻¹, after which a declining trend was observed at 16 g kg⁻¹ under both pre- and post-challenge conditions. However, no significant differences ($P>0.05$) were observed in PCV, MCV, or MCH between the experimental groups and the control group.

Serum biochemical assessment for liver enzymes:

The serum biochemical parameters of *O. niloticus* fed diets supplemented with different levels of *O. gratissimum* leaf extract before and after challenge with *V. parahaemolyticus* are presented in Figure 3. ALT, AST, and glucose levels were significantly higher ($P<0.05$) in the control group compared with fish fed *Ocimum*-supplemented diets under both pre- and post-challenge conditions. In addition, the control group exhibited a significant increase ($P<0.05$) in ALT and glucose levels following challenge with *V. parahaemolyticus*, indicating increased physiological stress and possible hepatic damage. In contrast, fish receiving diets supplemented with

Ocimum leaf extract maintained lower ALT, AST, and glucose values throughout the experimental period.

ALP activity and albumin concentration were significantly lower ($P<0.05$) in the control group compared with fish fed *Ocimum*-supplemented diets. Total protein levels were significantly enhanced in the supplemented groups, with the highest values recorded in fish fed 16 g kg⁻¹ *Ocimum* leaf extract during the pre-challenge period (3.93 ± 0.52 g dL⁻¹) and in fish fed 8 g kg⁻¹ during the post-challenge period (4.57 ± 0.26 g dL⁻¹). In contrast, the lowest total protein values were observed in the control group, with 1.24 ± 0.10 g dL⁻¹ and 0.70 ± 0.35 g dL⁻¹ recorded under pre- and post-challenge conditions, respectively.

Similarly, globulin concentrations were significantly higher ($P<0.05$) in fish fed *Ocimum*-supplemented diets compared with the control group. The highest globulin values were recorded in fish fed 8 g kg⁻¹ *Ocimum* leaf extract, reaching 2.87 ± 0.09 g dL⁻¹ and 2.99 ± 0.43 g dL⁻¹ under pre- and post-challenge conditions, respectively, whereas the control group exhibited the lowest values (0.79 ± 0.10 and 0.77 ± 0.30 g dL⁻¹, respectively). Albumin concentrations followed a similar trend, with the highest values observed in fish fed 8 g kg⁻¹ *Ocimum* leaf extract during the pre-challenge period (1.98 ± 0.16 g dL⁻¹) and 4 g kg⁻¹ during the post-challenge period (1.70 ± 0.29 g dL⁻¹). Conversely, the control group consistently recorded the lowest albumin values, with 0.55 ± 0.17 and 0.35 ± 0.14 g dL⁻¹ under pre- and post-challenge conditions, respectively.

Oxidative stress biomarkers: The oxidative stress biomarkers in the liver of *O. niloticus* fed diets supplemented with different levels of *O. gratissimum* leaf extract before and after challenge with *V. parahaemolyticus* are presented in Figure 4. SOD activity was significantly higher ($P<0.05$) in fish fed *Ocimum*-supplemented diets compared with the control group under both pre- and post-challenge conditions. The highest SOD activities were recorded in fish fed 8 g kg⁻¹ *Ocimum* leaf extract, with values




Figure 3. Serum biochemical parameters. (a) Alanine aminotransferase, (b) aspartate aminotransferase, (c) alkaline phosphatase, (d) total protein, (e) albumin, (f) globulin, and (g) glucose of *Oreochromis niloticus* fed a diet with different levels of *Ocimum gratissimum* leaf extract pre- and post-challenged with *Vibrio parahaemolyticus*. Values are mean $\pm$ SE; bars with different letters (a, b, c) differ significantly at $P < 0.05$. Upper case letters (A, B) indicated a significant difference at $P < 0.05$ between pre- and post-challenging with *V. parahaemolyticus*.

of 54.92 ± 2.59 IU mL⁻¹ and 55.27 ± 2.29 IU mL⁻¹ before and after challenge, respectively, whereas the control group exhibited the lowest values (33.30 ± 2.03 and 35.15 ± 1.29 IU mL⁻¹, respectively).

A similar pattern was observed for CAT activity, which increased significantly in fish receiving *Ocimum*-supplemented diets. The highest CAT activities were observed in fish fed 16 g kg⁻¹

(17.46 ± 1.25 U mL⁻¹) and 8 g kg⁻¹ (15.90 ± 2.08 U mL⁻¹) *Ocimum* leaf extract during the pre- and post-challenge conditions, respectively. In contrast, malondialdehyde (MDA) levels, an indicator of lipid peroxidation and oxidative damage, were significantly reduced ($P < 0.05$) in fish fed *Ocimum*-supplemented diets compared with the control group. The lowest MDA concentrations were recorded in fish fed 16 g




Figure 4. Oxidative stress biomarkers in the liver. (a) Superoxide dismutase, (b) catalase, (c) malondialdehyde, and (d) reduced glutathione of *Oreochromis niloticus* fed a diet with different levels of *Ocimum gratissimum* leaf extract pre- and post-challenged with *Vibrio parahaemolyticus*. Values are mean $\pm$ SE; bars with different letters (a, b, and c) differ significantly at $P<0.05$. Upper case letters (A and B) indicated a significant difference at $P<0.05$ between pre- and post-challenged with *V. parahaemolyticus*.

kg⁻¹ (27.33 $\pm$ 0.73 nmol mL⁻¹) and 8 g kg⁻¹ (38.80 $\pm$ 2.52 nmol mL⁻¹) *Ocimum* leaf extract under pre- and post-challenge conditions, respectively.

Reduced glutathione (GSH) levels were also significantly elevated ($P<0.05$) in fish fed *Ocimum* leaf extract, with the highest values observed in the 8 and 16 g kg⁻¹ dietary groups under both pre- and post-challenge conditions. Conversely, a significant reduction ($P<0.05$) in GSH levels was observed in the control group following *V. parahaemolyticus* challenge, indicating increased oxidative stress in untreated fish.

Immune parameters: Significant differences ($P<0.05$) in lysozyme activity were observed in the

spleen, skin, and kidney tissues of *O. niloticus* fed diets supplemented with different levels of *O. gratissimum* leaf extract before and after challenge with *V. parahaemolyticus* (Fig. 5). Among the examined tissues, lysozyme activity was highest in the kidney, followed by the spleen, and lowest in the skin. In all three organs, lysozyme activity increased progressively with increasing dietary inclusion levels of *Ocimum* leaf extract. Following bacterial challenge, the highest lysozyme activity was observed in fish fed 8 g kg⁻¹ *Ocimum* leaf extract in the kidney, whereas peak values in the spleen and skin were recorded in fish fed 16 g kg⁻¹. Overall, fish fed *Ocimum*-supplemented diets exhibited significantly higher




Figure 5. Lysozyme activity of (a) the spleen, (b) skin, and (c) kidney of *Oreochromis niloticus* fed a diet with different levels of *Ocimum gratissimum* leaf extract pre- and post-challenge with *Vibrio parahaemolyticus*. Values are the mean $\pm$ SE; bars with different letters (a, b, and c) differ significantly at $P < 0.05$. Uppercase letters (A and B) indicated a significant difference at $P < 0.05$ between pre- and post-challenge with *V. parahaemolyticus*.




Figure 6. Immunoglobulin (IgM) activity (a) in the spleen, (b) in the skin, and (c) in the kidney of *Oreochromis niloticus* fed a diet with different levels of *Ocimum gratissimum* Leaf Extract pre- and post-challenge with *Vibrio parahaemolyticus*. Values are the mean $\pm$ SE; bars with different letters (a, b, and c) differ significantly at $P < 0.05$. Uppercase letters (A and B) indicated a significant difference at $P < 0.05$ between pre- and post-challenge with *V. parahaemolyticus*.

($P < 0.05$) lysozyme activity in all examined tissues compared with the control group under both pre- and post-challenge conditions. However, no significant differences ($P > 0.05$) in lysozyme activity were detected between pre- and post-challenge conditions within the same dietary groups, irrespective of tissue type.

The immunoglobulin M (IgM) levels in the spleen, skin, and kidney are presented in Figure 6. Fish fed diets supplemented with *Ocimum* leaf extract showed significantly higher ($P < 0.05$) IgM levels compared with the control group under both pre- and post-challenge conditions. Among the analyzed tissues, IgM activity was highest in the skin, followed by the




Figure 7. Interleukine-1 β (IL-1 β) activity in (a) spleen, (b) skin, and (c) kidney of *Oreochromis niloticus* fed a diet with different levels of *Ocimum gratissimum* leaf extract pre- and post-challenge with *Vibrio parahaemolyticus*. Values are the mean $\pm$ SE; bars with different letters (a, b, and c) differ significantly at $P < 0.05$. Uppercase letters (A and B) indicated a significant difference at $P < 0.05$ between pre- and post-challenge with *V. parahaemolyticus*




Figure 8. Percentage survival of *Oreochromis niloticus* fed different levels of *Ocimum gratissimum* leaf extract in the diet for 90 days and challenged with *Vibrio parahaemolyticus*. Values are mean $\pm$ SE; bars with different letters (a, b, and c) differ significantly at $P < 0.05$. Uppercase letters (A and B) indicated a significant difference at $P < 0.05$ between pre- and post-challenge with *V. parahaemolyticus*




Figure 9. Skin lesion morbidity (%) of *Oreochromis niloticus* fed different levels of *Ocimum gratissimum* leaf extract challenged with *Vibrio parahaemolyticus*. Values are the mean $\pm$ SE; bars with different letters (a, b, and c) differ significantly at $P < 0.05$

kidney, and lowest in the spleen. Prior to challenge, the highest IgM levels were recorded in fish fed 4 g kg⁻¹ *Ocimum* leaf extract in the skin, 16 g kg⁻¹ in the spleen, and 8 g kg⁻¹ in the kidney. Following challenge with *V. parahaemolyticus*, IgM levels increased further in all tissues, reaching peak values in fish fed 16 g kg⁻¹ *Ocimum* leaf extract. Significant differences ($P < 0.05$) in IgM levels were observed between the treatment groups and the control following bacterial challenge.

The interleukin-1 β (IL-1 β) levels in the spleen, skin, and kidney tissues of *O. niloticus* are shown in Figure 7. The control group exhibited significantly higher IL-1 β levels ($P < 0.05$) than fish fed *Ocimum*-supplemented diets under both pre- and post-challenge conditions. Among the examined tissues, IL-1 β levels were highest in the skin, followed by the spleen, and lowest in the kidney. Following challenge with *V. parahaemolyticus*, a significant increase ($P < 0.05$) in IL-1 β levels was observed in all tissues of the control group. Fish fed 2 g kg⁻¹ *Ocimum* leaf extract showed a significant increase ($P < 0.05$) in IL-1 β levels in the spleen and skin after challenge, whereas fish fed




Figure 10. Skin lesion morbidity of *Oreochromis niloticus* fed with (a) 0, (b) 2, (c) 4, (d) 8, and (e) 16 g/kg of *Ocimum gratissimum* leaf extract for 90 days, respectively. a, b, c, d, and e- showing normal skin.

4 g kg⁻¹ exhibited a significant increase only in the skin. In contrast, fish fed 8 and 16 g kg⁻¹ *Ocimum* leaf extract showed no significant differences ($P>0.05$) in IL-1 β levels between pre- and post-challenge conditions, indicating a potential modulatory effect of the extract on inflammatory responses.

Challenge test: No significant differences ($P>0.05$) in survival rate were observed among the experimental groups and the control group following the 90-day feeding trial prior to bacterial challenge (Fig. 8). However, after challenge with *V. parahaemolyticus*,

fish fed diets supplemented with *O. gratissimum* leaf extract exhibited higher survival rates compared with the control group. The highest post-challenge survival rate (80%) was recorded in fish fed diets containing 8 and 16 g kg⁻¹ *Ocimum* leaf extract, whereas the lowest survival rate (30%) was observed in the control group receiving no dietary supplementation.

The severity of skin lesion morbidity following bacterial challenge is presented in Figures 9-11. Fish in the control group exhibited the highest percentage of skin lesion morbidity, whereas fish fed *Ocimum*-




Figure 11. Post-challenge skin lesion morbidity of *Oreochromis niloticus* fed a diet with (a) 0, (b) 2, (c) 4, (d) 8 and (e) 16 g/kg diet *Ocimum gratissimum* leaf extract, respectively. ai- showing severe skin necrosis (SSU) and fin necrosis (FN); showing loss of scale (LS); bi- showing skin haemorrhage (SH); bii- showing skin necrosis (SN); ci- showing skin necrosis (SN); cii- showing skin haemorrhage (SH); di- showing skin necrosis (SN); dii- showing fin necrosis (FN) and ei- showing skin necrosis (SU); eii- showing skin haemorrhage (SH)

supplemented diets showed markedly reduced lesion severity. The highest skin lesion morbidity (80%) was recorded in the control group, while the lowest value (15%) was observed in fish fed 8 g kg⁻¹ *Ocimum* leaf extract, indicating improved resistance against *V. parahaemolyticus* infection in the supplemented groups.

Discussions

The present study demonstrated that dietary supplementation with *O. gratissimum* leaf extract

improved the hematological status and immune responsiveness of *O. niloticus*, particularly at the inclusion level of 8 g kg⁻¹. Fish fed *Ocimum*-supplemented diets exhibited significantly higher WBC, RBC, HGB, and PCV values compared with the control group under both pre- and post-challenge conditions. In contrast, the control group showed a marked reduction in HGB concentration following *V. parahaemolyticus* challenge, indicating greater physiological stress and susceptibility to infection. However, no significant differences were observed in

MCV and MCH among the experimental groups. The increased WBC counts observed in fish fed *Ocimum*-supplemented diets suggest enhanced nonspecific immune responses and improved defense mechanisms against pathogenic infection. Similar findings have been reported in *Clarias gariepinus*, rainbow trout (*Oncorhynchus mykiss*), and common carp (*Cyprinus carpio*) fed diets supplemented with phytogetic additives, including *Carica papaya* (paw-paw) aqueous leaf extract (Oparaku et al., 2024), *Plantago major* leaf extract (Khanzadeh et al., 2026), and papaya leaf extract (Giri et al., 2026).

The elevation of WBC levels after bacterial challenge further indicates immune system activation in response to *V. parahaemolyticus* infection. Likewise, the higher RBC and HGB values observed in the supplemented groups indicate improved erythropoietic activity and oxygen-carrying capacity, reflecting better physiological condition and stress tolerance. Comparable improvements in RBC and HGB levels have also been reported in fish fed medicinal plant-based diets containing *Terminalia arjuna* bark (Nandi et al., 2026), spirulina and black cumin (Ayesha et al., 2026), and *Hunteria umbellata* (Jimoh et al., 2026). The reduction in HGB observed in the control group after bacterial challenge may be associated with hemolysis, impaired erythropoiesis, or infection-induced stress, whereas fish fed *Ocimum*-supplemented diets maintained relatively stable HGB levels, suggesting a protective effect against hematological disturbances. The lack of significant changes in MCV and MCH indicates that *Ocimum* leaf extract mainly influenced blood cell production rather than erythrocyte morphology. The beneficial effects observed in the present study are likely attributable to the rich phytochemical composition of *O. gratissimum*, particularly flavonoids, phenolics, thymol, and eugenol, which possess antioxidant and immunostimulatory properties that can enhance hematopoiesis, reduce oxidative stress, and stimulate immune-related pathways.

The present study demonstrated that dietary supplementation with *O. gratissimum* leaf extract

improved the serum biochemical profile and hepatic health status of *O. niloticus*, particularly following *V. parahaemolyticus* challenge. Fish fed *Ocimum*-supplemented diets exhibited significantly lower ALT, AST, and glucose levels, whereas ALP, total protein, globulin, and albumin concentrations were significantly elevated compared with the control group. In contrast, the marked increase in ALT and glucose levels observed in the control group after bacterial challenge suggests greater physiological stress and hepatic impairment. Similar improvements in serum biochemical parameters have been reported in fish fed phytogetic additives, including *C. papaya* aqueous leaf extract (Oparaku et al., 2024), *P. major* leaf extract (Khanzadeh et al., 2026), papaya leaf extract (Giri et al., 2026), and fish mint (*Houttuynia cordata* Thunb.) (Yu et al., 2026). Elevated ALT and AST activities are commonly recognized as indicators of hepatocellular damage and physiological stress; thus, the lower enzyme activities observed in the supplemented groups suggest hepatoprotective effects of *O. gratissimum* (Refaey et al., 2023). Likewise, the reduced glucose levels in fish fed *Ocimum*-supplemented diets may reflect improved stress tolerance and metabolic regulation during bacterial infection. The increased concentrations of total protein, globulin, and albumin further indicate enhanced protein synthesis, immune competence, and overall physiological status (Reshi et al., 2023). These beneficial effects are likely attributed to the rich phytochemical composition of *O. gratissimum*, particularly flavonoids, phenolics, thymol, and eugenol, which possess antioxidant, antimicrobial, and hepatoprotective activities capable of mitigating oxidative damage and improving liver function in challenged fish (Silva et al., 2022; Ugbogu et al., 2021).

The present study showed that dietary supplementation with *O. gratissimum* leaf extract enhanced the antioxidant defense system of *O. niloticus*, as evidenced by increased SOD, catalase CAT, and reduced GSH activities, together with reduced MDA levels under both pre- and post-

challenge conditions. The strongest antioxidant responses were generally observed in fish fed 8 g kg⁻¹ *Ocimum* leaf extract, whereas the control group exhibited lower antioxidant enzyme activities and higher oxidative stress indicators following *V. parahaemolyticus* challenge. Similar improvements in antioxidant biomarkers have been reported in fish fed phytogetic additives such as *Ocimum* species (Abdel-Tawwab et al., 2018), *Perilla frutescens*, *Illicium verum*, and *Artemisia annua* extracts (Yue et al., 2024), Scots pine (*Pinus sylvestris*) essential oil (Yousefi et al., 2025), and liposome encapsulating pine bark extract (Ibrahim et al., 2025). Antioxidant enzymes, including SOD and CAT, play essential roles in protecting cells against reactive oxygen species (ROS) and oxidative damage (Hoseinifar et al., 2020; Khan et al., 2025). Therefore, the elevated SOD and CAT activities observed in the supplemented groups indicate enhanced antioxidant defense capacity and improved protection against infection-induced oxidative stress (Jomova et al., 2024). Likewise, the higher GSH levels further support the antioxidative potential of *O. gratissimum*, whereas the reduction in GSH observed in the control group after bacterial challenge suggests depletion of antioxidant reserves due to excessive oxidative stress. MDA, a marker of lipid peroxidation and cellular membrane damage (Rizzo, 2024), was significantly reduced in fish fed *Ocimum*-supplemented diets, indicating lower oxidative damage and improved membrane stability. These beneficial effects are likely associated with the rich phytochemical composition of *O. gratissimum*, particularly flavonoids, phenolics, thymol, and eugenol, which possess strong antioxidant and free radical scavenging activities capable of enhancing endogenous antioxidant defenses and mitigating oxidative damage during bacterial infection (Akinmoladun et al., 2007; Mahapatra and Roy, 2014; Ballester et al., 2025).

The results of the current work revealed that dietary supplementation with *O. gratissimum* leaf extract enhanced the immune responses of *O. niloticus*, as evidenced by increased lysozyme and IgM activities

and reduced IL-1 β levels under both pre- and post-challenge conditions. Fish fed *Ocimum*-supplemented diets exhibited significantly higher lysozyme and IgM levels in the spleen, skin, and kidney compared with the control group, whereas IL-1 β levels were significantly lower, particularly at 8 and 16 g kg⁻¹ inclusion levels. These findings suggest that dietary *O. gratissimum* enhanced both innate and adaptive immune responses and modulated excessive inflammatory responses induced by *V. parahaemolyticus* infection. Similar immunostimulatory effects have been reported in Nile tilapia and other fish species fed phytogetic additives, including *Ocimum* species, garlic, ginger, and medicinal herb extracts (Abdel-Tawwab et al., 2018; Ergena et al., 2023; Omitoyin et al., 2019) and *Avicennia marina* leaf extract (El Basuini, 2025). Lysozyme is an important component of the innate immune system involved in bacterial cell wall degradation and pathogen elimination (Mokhtar et al., 2023). Therefore, the elevated lysozyme activity observed in the supplemented groups indicates enhanced nonspecific immune defense. Likewise, the increased IgM levels suggest improved humoral immunity and antibody production (Huang et al., 2025). In contrast, the elevated IL-1 β levels observed in the control group after bacterial challenge indicate stronger inflammatory responses associated with infection-induced stress and tissue damage. The lower IL-1 β levels recorded in fish fed *Ocimum*-supplemented diets suggest anti-inflammatory effects and better regulation of immune responses during infection (Mansour et al., 2023). These beneficial effects are likely associated with the rich phytochemical composition of *O. gratissimum*, particularly flavonoids, phenolics, thymol, and eugenol, which possess antimicrobial, antioxidant, immunostimulatory, and anti-inflammatory properties that enhance immune competence and disease resistance in Nile tilapia.

The challenge test demonstrated that dietary supplementation with *O. gratissimum* leaf extract enhanced *O. niloticus* resistance to

V. parahaemolyticus infection, as evidenced by higher post-challenge survival rates and lower skin lesion severity in the supplemented groups compared with the control group. Fish fed diets containing 8 and 16 g kg⁻¹ *Ocimum* leaf extract showed the highest survival rate (80%), whereas the control group exhibited the lowest survival (30%). In addition, skin lesion morbidity was markedly reduced in fish receiving *Ocimum*-supplemented diets, particularly at 8 g kg⁻¹ inclusion level. Similar improvements in disease resistance and survival have been reported in fish fed phytogenic additives such as *Gentiana Macrophylla* pall aqueous extract (Zhou et al., 2024), herbal extracts (Zhu et al., 2024), herbal extract blend (Zeng et al., 2025), papaya leaf extract (Giri et al., 2026), and α -mangostin-rich extract nanoemulsion (Srisaen et al., 2026). The lower survival and higher lesion severity observed in the control group may be associated with increased bacterial colonization, oxidative stress, and weakened immune responses following infection. In contrast, the improved survival and reduced lesion formation in the supplemented groups were likely related to enhanced hematological, antioxidant, and immune responses induced by *O. gratissimum*. The protective effects of *O. gratissimum* are likely attributable to its rich phytochemical composition, particularly flavonoids, phenolics, thymol, and eugenol, which possess antimicrobial, antioxidant, anti-inflammatory, and immunostimulatory properties that enhance disease resistance and physiological resilience in Nile tilapia.

Acknowledgement

The authors thank Kebbi State University of Science and Technology, Aliero, for PhD sponsorship and the Department of Biology for providing the facilities for the conduct of this research.

References

- Abdel-Tawwab M., Adeshina I., Jenyo-Oni A., Ajani E.K., Emikpe B.O. (2018). Growth, physiological, antioxidants, and immune response of African catfish, *Clarias gariepinus* (B.), to dietary clove basil, *Ocimum gratissimum*, leaf extract and its susceptibility to *Listeria monocytogenes* infection. Fish and Shellfish Immunology, 78: 346-354.
- Abeysinghe K.C., Weeraratne A.G.P.T., Colonne D.K.C., Bellanthudawa B.K.A. (2025). Sustainable development goals (SDGs), climate change, and the development of aquaculture and fisheries industries. Environmental Reviews, 33: 1-24.
- Adhikary U., Ahmmmed S.S., Sadia H.T., Sarkar R., Nasrin F., Ghosh A.K. (2026). Dietary inclusion of *Allium cepa* and *Allium sativum* extract improves growth performance, haematological and immunological indices, body composition and disease resistance of prawn (*Macrobrachium rosenbergii*) against *Vibrio parahaemolyticus* infection. Fish and Shellfish Immunology, 169: 111021.
- Aebi H. (1984). Oxidoreductases acting on groups other than CHOH: Catalase. Methods in Enzymology, 105: 121-125.
- Akanbi D.A., Ogboriga P.O., Adeniyi I.A., Lawal A.M., Olusola M.A., Oluyemi P.A., Onasanwo S.A. (2026). The neuroprotective effects of *Ocimum gratissimum*-supplemented diet on scopolamine-induced memory impairment in mice model of Alzheimer's Disease. Phytomedicine Plus, 6(2): 100979.
- Akinmoladun A., Ibukun E., Afor E., Obuotor E.M., Farombi O. (2007). Phytochemical constituent and antioxidant activity of extract from the leaves of *Ocimum gratissimum*. Scientific Research and Essay, 2: 163-166.
- Ayesha S.B., Islam M.J., Mimi M.S.A., Sheuly M.S.A., Auditi T.I., Hossain M.K., Sumon K.A., Shahjahan M., Alam M.S., Rashid H. (2026). Dietary spirulina and black cumin supplementation mitigate extreme warm temperature stress in Nile tilapia, *Oreochromis niloticus*. Aquaculture and Fisheries. <https://doi.org/10.1016/j.aaf.2026.02.005>
- Bai S.C., Hamidoghli A., Davies S.J., Wan A.H.L. (2025). Chapter 7 - Feed additives. In: V. Kumar (Ed.), Feed and Feeding for Fish and Shellfish. Academic Press. pp: 173-211.
- Ballester E.L.C., Costa W.G.d.S., do Nascimento Ferreira C.H., Retcheski M.C., Cazarolli L.H., Schwengber G.H., Mauerwerk M.T., Pinheiro C.G., Heinzmann B.M., Baldisserotto B., de Souza Valente C. (2025). Dietary *Ocimum gratissimum* essential oil improves the antioxidant status and maintains the performance of *Macrobrachium rosenbergii* juveniles. Applied

- Sciences, 15(5): 2745.
- Bandeira G., Pês T.S., Saccol E.M.H., Sutili F.J., Rossi W., Murari A.L., Heinzmann B.M., Pavanato M.A., de Vargas A.C., de L. Silva L., Baldisserotto B. (2017). Potential uses of *Ocimum gratissimum* and *Hesperozygis ringens* essential oils in aquaculture. *Industrial Crops and Products*, 97: 484-491.
- Beutler E., Duron O., Kelly B.M. (1963). Improved method for determination of blood glutathione. *Journal of Laboratory and Clinical Medicine*, 61: 882-8.
- Camberos-Solano M., Soto-Rodriguez S.A. (2026). Toxin Secretion Systems Used by Marine Pathogenic Bacteria: A Review. *Journal of Fish Diseases*, e70194.
- Chakroborty K., Lima R.A., Foyzul Hossain M., Rafiquzzaman S.M. (2025). Biobased functional feed additives in asian aquaculture: Trends, impacts, and future directions. *Animal Feed Science and Technology*, 116222.
- Chupani L., Niksirat H., Erasmus B., Piackova V., Zuskova E. (2026). Sustainable parasite management in aquaculture: Integrating productivity and environmental health. *Aquaculture*, 622: 744026.
- Dacie J., Lewis S. (1975). *Practical haematology*. London Churchill Livingstone.
- Egba S.I., Ugwu C.E., Ihediwa F.D. (2014). Changes In Cytokine Expression In Rats Administered Extracts of *Termitomyces robustus* And *Ocimum gratissimum*. *International Journal of Life Sciences Biotechnology and Pharma Research*, 3(4): 182.
- El Basuini M.F. (2025). Immunomodulatory, antioxidant, and growth-promoting properties of *Avicennia marina* leaf extract on Nile tilapia. *Scientific Reports*, 15(1): 43948.
- El-Sayed A.-F.M., Fitzsimmons K. (2023). From Africa to the world—The journey of Nile tilapia. *Reviews in Aquaculture*, 15(S1): 6-21.
- ELTarabili R.M., Elsayed M.E., Elghayaty H., El Kattawy Z.M. (2026). Emergence of Multidrug-Resistant *Vibrio parahaemolyticus* and *Vibrio alginolyticus* in the Nile Tilapia from Manzala Wetland. *Egyptian Journal of Aquatic Biology and Fisheries* 30(2): 2745-2757.
- Emam W., Lambert H., Brown C. (2025). The welfare of farmed Nile tilapia: a review. *Frontiers in Veterinary Science*, 12. <https://doi.org/10.3389/fvets.2025.1567984>
- Ergena A., Natarajan P., Bedewi Z. (2023). Determination of some haematological parameters and disease resistance capacity to *Aeromonas hydrophila* infection in the Nile tilapia, *Oreochromis niloticus* L. fed dietary supplementation of ginger (*Zingiber officinale*). *Egyptian Journal of Aquatic Biology and Fisheries*, 27(3): 551-568.
- Giri S.S., Jun J.W., Lee S.B., Joo S.J., Hwang M.H., Park J.H., Park D.S., Park E.J., Sukumaran V., Chang Park S. (2026). Effect of papaya leaf extract on growth performance, skin mucosal immune responses, liver enzymes, disease resistance, and tight junctions of *Cyprinus carpio*. *Fish and Shellfish Immunology*, 168: 110916.
- Hassan H.U., Ali A., Al Sulivany B.S.A., Bilal M., Kanwal R., Raza M.A., Arslan A., Ijaz M.Z., Kabir M., Khan M.R., Rasdi N.W., Arai T. (2025). Investigation of the effects of phytogenic dietary additives on growth performance, nutrient utilization, economic efficiency and health of *Pangasius hypophthalmus*: implications for sustainable aquaculture development. *Scientific Reports*, 15(1): 22661.
- Hoseinifar S.H., Yousefi S., Van Doan H., Ashouri G., Gioacchini G., Maradonna F., Carnevali O. (2020). Oxidative stress and antioxidant defense in fish: The implications of probiotic, prebiotic, and synbiotics. *Reviews in Fisheries Science and Aquaculture*, 29(2): 198-217.
- Hossain M.M.M., Rojoni S.A., Ferdous M.J., Rahman N., Rubayea U., Bristy S.A., Mojumdar S., Sen B.K., Lopa N.J., Ghosh R.R., Suhag N.I., Rahman M., Hasan M.M., Rahman M.A. (2026). Plant-mediated synthesis: Transforming traditional Bangladeshi medicinal plants into immunomodulatory nanoparticles for enhanced shrimp immunity and pathogen control. *Microbial Pathogenesis*, 210: 108144.
- Hu Y., Liang Y., Han L., Hu D., Li M., Zhou Z., Hu Y., Wang X., Yin Y. (2025). Phytogenic diet enhances the growth and disease resistance of the black-spotted frog, *Pelophylax nigromaculatus*, by stabilizing intestinal homeostasis and boosting innate immunity. *Fish and Shellfish Immunology*, 165: 110553.
- Huang Z., Liu N., Xue M., Xu C., Fan Y., Meng Y., Jiang N., Li Y., Liu W., He Y., Zhou Y. (2025). Immunoglobulin M response in largemouth bass (*Micropterus salmoides*) following ranavirus infection. *Frontiers in Immunology*, 16. <https://doi.org/10.3389/fimmu.2025.1515684>
- Ibrahim D., Shahin S.E., Elnahriry S.S., El-Badry S.M.,

- Eltarabili R.M., Elazab S.T., Ismail T.A., Abd El-Hamid M.I. (2025). Liposome encapsulating pine bark extract in Nile tilapia: Targeting interrelated immune and antioxidant defense to combat coinfection with *Aeromonas hydrophila* and *Enterococcus faecalis*. *Fish and Shellfish Immunology*, 156: 110031.
- Ivanova S., Sukhikh S., Popov A., Shishko O., Nikonov I., Kapitonova E., Krol O., Larina V., Noskova S., Babich O. (2024). Medicinal plants: A source of phytobiotics for the feed additives. *Journal of Agriculture and Food Research*, 16: 101172.
- Jimoh W.A., Ayeloja A.A., Olonikadi H.F., Abdulwaheed A.A., Yusuff H.A., Alli F.E., Ajayi J.A., Taiwo A.D., Fawole F.J. (2026). Growth performance and physiological responses of hybrid catfish (*Clarias gariepinus* ♀ × *Heterobranchius bidorsalis* ♂) to dietary *Hunteria umbellata* supplementation. *Animal Feed Science and Technology*, 340: 116798.
- Jomova K., Alomar S.Y., Alwasel S.H., Nepovimova E., Kuca K., Valko M. (2024). Several lines of antioxidant defense against oxidative stress: antioxidant enzymes, nanomaterials with multiple enzyme-mimicking activities, and low-molecular-weight antioxidants. *Archives of Toxicology*, 98(5): 1323-1367.
- Kamelnia E., Mohebbati R., Kamelnia R., El-Seedi H.R., Boskabady M.H. (2023). Anti-inflammatory, immunomodulatory and anti-oxidant effects of *Ocimum basilicum* L. and its main constituents: A review. *Iranian Journal of Basic Medical Sciences*, 26(6): 617-627.
- Khan M., Mushtaq M., Usman M., Rahman M.A.U., Quan G. (2025). Oxidative stress-induced cytotoxicity and the role of dietary antioxidants in farm animals: A review. *Advances in Redox Research*, 16: 100138.
- Khanzadeh M., Hoseinifar S.H., Mazandarani M., Safari R., Jafari V., Van Doan H. (2026). Effects of *Plantago major* leaf extract on growth, serum and mucosal immunity responses, and skin-related gene expression in triploid juvenile rainbow trout (*Oncorhynchus mykiss*). *Aquaculture Reports*, 46: 103375.
- Koley A., Gupta N., Banerjee S., Patchaiyappan A., Bagdi T., Ruchi R.D., Debnath S., Chaudhury S., Muthusamy A., Hazra A.K., Balachandran S. (2025). Sustainable bio-valorisation of water hyacinth for enhancing rural livelihood and achieving sustainable development goals. *Bioresource Technology Reports*, 32: 102352.
- Leite dos Santos C.A., Silva Pereira R.L.d., Pereira-de-Morais L., Santos G.J.G., Coutinho H.D.M., Assis Bezerra da Cunha F., Rodrigues F.F.G., Sousa Júnior D.L.D., Silva J.C. (2025). Antibacterial activity of ethanolic extract of *Ocimum gratissimum* L. (BASIL) and its toxicity against *Drosophila melanogaster*. *Pharmacological Research - Natural Products*, 7: 100212.
- Li M., Jiang H., Shi K., Huang H., Wang X., Li F., Pan K., Yang Y., Zhang R. (2026). Efficient elimination of highly lethal *Vibrio parahaemolyticus* in shrimp aquaculture by a vibriophage. *Fish and Shellfish Immunology*, 169: 111037.
- Liboureau P., Mols-Mortensen A., Laznya A., Pampanin D.M. (2026). Seaweed aquaculture as a sustainable source of functional foods. *Future Foods*, 13: 100876.
- Liu H.-X., Zhou X.-Q., Jiang W.-D., Wu P., Liu Y., Zeng Y.-Y., Jiang J., Kuang S.-Y., Tang L., Feng L. (2018). Optimal α -lipoic acid strengthen immunity of young grass carp (*Ctenopharyngodon idella*) by enhancing immune function of head kidney, spleen and skin. *Fish and Shellfish Immunology*, 80: 600-617.
- Maezono M., Nielsen R., Buchmann K., Nielsen M. (2025). The current state of knowledge of the economic impact of diseases in global aquaculture. *Reviews in Aquaculture*, 17(3): e70039.
- Mahapatra S.K., Roy S. (2014). Phytopharmacological approach of free radical scavenging and anti-oxidative potential of eugenol and *Ocimum gratissimum* Linn. *Asian Pacific Journal of Tropical Medicine*, 7(s1): S391-397.
- Manchanayake T., Salleh A., Amal M.N.A., Yasin I.S.M., Zamri-Saad M. (2023). Pathology and pathogenesis of *Vibrio* infection in fish: A review. *Aquaculture Reports*, 28: 101459.
- Mansour A.T., Diab A.M., Khalil R.H., Eldessouki E.A., El-Sabbagh N., Elsamannoudy S.I., Younis N.A., (2023). Physiological and immunological responses of Nile tilapia fed dietary supplementation of sweet basil ethanolic and aqueous extracts. *Frontiers in Marine Science*, 9. <https://doi.org/10.3389/fmars.2022.1064455>
- Mokhtar D.M., Zacccone G., Alesci A., Kuciel M., Hussein M.T., Sayed R.K.A. (2023). Main Components of Fish Immunity: An Overview of the Fish Immune System. *Fishes*, 8(2): 93.
- Nandi S.K., Hridoy M.A.A.M., Sarkar S., Abdul Kari Z., Siddik M.A.B., Hasan A.M.N., Eissa E.-S.H., Van

- Doan H., Reduan M.F.H., Kabir M.A. (2026). Zootechnical performance, disease resistance and economic efficiency of climbing perch, *Anabas testudineus* fed Terminalia arjuna bark-supplemented diets. Aquaculture Reports, 46: 103328.
- Ndaro M.E., Mzula A., Funga A., Rumisha C. (2026). Economic impact of disease outbreaks on cage fish farming in Tanzania's Lake Victoria Basin. Aquaculture, Fish and Fisheries, 6(2): e70243.
- Nishikimi M., Rao N.A., Yagi K. (1972). The occurrence of superoxide anion in the reaction of reduced phenazine methosulfate and molecular oxygen. Biochemical and Biophysical Research Communications, 46(2): 849-854.
- Nordhagen S., Beal T., Herrero M., Mane E., Neufeld L.M., Sokourenko K. (2026). Livelihood implications of dietary shifts to animal-source food alternatives: a scoping review. The Lancet Planetary Health, 10(2): 101424.
- Odeniyi O.A., Adekunle Y.A., Adebayo P.A., Adedeji T.S. (2025). Phytochemical compositions, *in vitro* antimicrobial activity, and molecular docking analysis of essential oils of *Ocimum gratissimum* and *Citrus sinensis* against selected clinical isolates. Next Research, 2(4): 101084.
- Ohkawa H., Ohishi N., Yagi K. (1979). Assay for lipid peroxides in animal tissues by thiobarbituric acid reaction. Analytical Biochemistry, 95(2): 351-358.
- Omitoyin B.O., Ajani E.K., Orisasona O., Bassey H.E., Kareem K.O., Osho F.E. (2019). Effect of guava *Psidium guajava* (L.) aqueous extract diet on growth performance, intestinal morphology, immune response and survival of *Oreochromis niloticus* challenged with *Aeromonas hydrophila*. Aquaculture Research, 50(7): 1851-1861.
- Oparaku N.F., Ukwueze M., Nwosu C.G., Andong F., Echiude D., Okwuonu E., Ezenwaji N. (2024). Effect of *Carica papaya* (paw-paw) aqueous leaf extract on the growth, haematological parameters and the liver enzymes of *Clarias gariepinus* juveniles. Aquaculture Reports, 34: 101874.
- Paredes-Trujillo A., Mendoza-Carranza M. (2022). A systematic review and meta-analysis of the relationship between farm management, water quality and pathogen outbreaks in tilapia culture. Journal of Fish Diseases, 45(10): 1529-1548.
- Petrusic M.I., McLean S.K., Palombo E.A. (2025). Characterisation of newly identified vibriophages and their potential application in the biocontrol of *Vibrio parahaemolyticus* to enhance safety in aquaculture. Virus Research, 362: 199664.
- Pudota B.A., Tambireddy N., Chennu R., Chethurajupalli L., Shaik H., Nadella R.K., Chatterjee N.S., Paturi A.P. (2025). Exploring the perspectives of phytobiotics and their role in aquaculture: Present status and future trends. The Microbe, 8: 100496.
- Refaey M.M., Mehri A.I., El-Komy M.M., Zenhom O.A., Mansour A.T. (2023). Chronic cold-stress induced histopathological changes, oxidative stress, and alterations in liver functions and nutrient composition of hybrid red tilapia and the potential protection of unsaturated fatty acids. Frontiers in Marine Science, 10. <https://doi.org/10.3389/fmars.2023.1148978>
- Reshi Q.M., Ahmed I., Al-Anazi K.M., Farah M.A. (2023). Indexing hematological and serum biochemical reference intervals of Himalayan snow trout, *Schizothorax esocinus* to instrument in health assessment. Frontiers in Physiology, 14: 989442.
- Rizzo M. (2024). Measurement of malondialdehyde as a biomarker of lipid oxidation in fish. American Journal of Analytical Chemistry, 15: 303-332.
- Sagar A., Yash M., Shalini T., Vinay Kumar R. (2025). The healing potential of *Ocimum gratissimum*: An integrative review of its ethnomedicinal uses, phytochemistry, and pharmacological relevance. International Journal of Pharmaceutical Research and Development, 7(2): 454-469
- Salau A.K., Yakubu M.T., Oladiji A.T. (2013). Cytotoxic activity of aqueous extracts of *Anogeissus leiocarpus* and *Terminalia avicennioides* root barks against Ehrlich Ascites Carcinoma cells. Indian Journal of Pharmacology, 45(4): 381-385.
- Sanches-Fernandes G.M.M., Sá-Correia I., Costa R. (2022). Vibriosis outbreaks in aquaculture: Addressing Environmental and public health concerns and preventive therapies using gilthead seabream farming as a model system. Frontiers in Microbiology, 13: 904815.
- Sharma S., Salaria D., Rolta R., Dubey R.C. (2026). Antioxidant and inhibition of protein denaturation potential of essential oils of *Ocimum tenuiflorum* and *Ocimum gratissimum*: In vitro evaluation and In-silico PASS analysis. Pharmacological Research - Natural Products, 10: 100474.
- Silva J.C., Silva Pereira R.L., Sampaio de Freitas T., Rocha

- J.E., Macedo N.S., de Fatima Alves Nonato C., Linhares M.L., Arruda Tavares D.S., Bezerra da Cunha F.A., Melo Coutinho H.D., Gonçalo de Lima S., Pereira-Junior F.N., Araújo Maia F.P., Pita Neto I.C., Galvão Rodrigues F.F., Garcia Santos G.J. (2022). Evaluation of antibacterial and toxicological activities of essential oil of *Ocimum gratissimum* L. and its major constituent eugenol. *Food Bioscience*, 50: 102128.
- Soares M.P., Maraschi A.C., de Angelis C.F., Queiroz S.C.N., Jonsson C.M., de Campos C.M., Fernandes M.N., Leite C.A.C., Rantin F.T., Sampaio F.G. (2026). Boosting growth, health, and productivity of Nile tilapia by supplementing *Artemisia annua* under tropical net cage farming conditions. *Aquaculture Reports*, 48: 103514.
- Srisaen W., Sutthi N., Rinthong P.-O., Chaiyasing R., Prisingkorn W., Kersanté P., Nontasan S., Wangkahart E. (2026). Synergistic effects of α -mangostin-rich extract nanoemulsion and a natural free amino acid mixture on growth performance, immune function, intestinal microbiota, and disease resistance in Nile tilapia (*Oreochromis niloticus*). *Fish and Shellfish Immunology*, 168: 111009.
- Ugbogu O.C., Emmanuel O., Agi G.O., Ibe C., Ekweogu C.N., Ude V.C., Uche M.E., Nnanna R.O., Ugbogu, E.A., (2021). A review on the traditional uses, phytochemistry, and pharmacological activities of clove basil (*Ocimum gratissimum* L.). *Heliyon*, 7(11).
- Vilhena Batista D.V., Martins Brasil E., Brum A., Laterça Martins M., Pedreira Mourinho J.L., Owatari M.S. (2026). Edwardsiellosis in tilapia farming: Impacts, challenges, and innovations for sustainable control. *Aquaculture and Fisheries*. <https://doi.org/10.1016/j.aaf.2026.02.006>
- Wang J., Deng L., Chen M., Che Y., Li L., Zhu L., Chen G., Feng T. (2024). Phytogetic feed additives as natural antibiotic alternatives in animal health and production: A review of the literature of the last decade. *Animal Nutrition*, 17: 244-264.
- Xi X., Lin M., Huang X., Yan G., Luo C., Chen R., Liu G., Wang P., Zhang Y. (2026). Effects of dietary *Astragalus membranaceus* extract and fucoidan on growth performance, hepatic resilience, antioxidant defense and gut homeostasis of largemouth bass (*Micropterus salmoides*). *Aquaculture Reports*, 48: 103510.
- Yousefi M., Adineh H., Kulikov E.V., Karamyan A.S., Rudenko P.A., Sachivkina N., Hoseini S.M. (2025). Effects of dietary Scots pine (*Pinus sylvestris*) essential oil on growth performance, digestive enzymes, immunological and antioxidant responses to *Aeromonas hydrophila* infection in rainbow trout (*Oncorhynchus mykiss*). *Fish and Shellfish Immunology*, 167: 110899.
- Yu M., Chen B., Zou Z., Zhu J., Li D., Yu J., Yang H., Xiao W. (2026). Integrating fish mint (*Houttuynia cordata* Thunb.) floating beds and feed additives to improve health and disease resistance in Nile tilapia. *Animal Feed Science and Technology* 335, 116677.
- Yue R., Dong W., Feng Z., Jin T., Wang W., He Y., Chen Y., Lin S. (2024). Effects of three tested medicinal plant extracts on growth, immune function and microflora in juvenile largemouth bass (*Micropterus salmoides*). *Aquaculture Reports*, 36: 102075.
- Zaher H.A., Nofal M.I., Hendam B.M., Elshaer M.M., Alothaim A.S., Eraqi M.M. (2021). Prevalence and antibiogram of *Vibrio parahaemolyticus* and *Aeromonas hydrophila* in the flesh of Nile tilapia, with special reference to their virulence genes detected using the multiplex PCR technique. *Antibiotics* (Basel), 10(6). <https://doi.org/10.3390/antibiotics10060654>
- Zeng Z.Y., Yu D.D., Shen Y.T., He X.H., Yang S., Fei H. (2025). Influence of herbal extract blend supplementation on growth performance, liver and intestinal health, intestinal transcriptome, and resistance to *Aeromonas veronii* in *Micropterus salmoides* with optimized feeding intervals. *Animal Feed Science and Technology*, 328: 116467.
- Zhou Y., Liu L., Song C., Hu Y., Chen J. (2024). The potent potential of *Gentiana macrophylla* pall aqueous extract in protecting largemouth bass against *Micropterus Salmoides* Rhabdovirus: A promising application of herbal crude extract in the fight against MSRV. *Aquaculture*, 578: 740069.
- Zhu C.B., Shen Y.T., Ren C.H., Yang S., Fei H. (2024). A novel formula of herbal extracts regulates growth performance, antioxidant capacity, intestinal microbiota and resistance against *Aeromonas veronii* in largemouth bass (*Micropterus salmoides*). *Aquaculture*, 583: 740614.